

水産総合研究センター研究報告第11号で掲載した2002年度論文等の発表リストにおいて記載漏れがありましたので、以下のとおり、修正致します。

水産総合研究センターにおける2002年度論文等の発表リスト。なお、\* 2001年度論文等で7号に記載漏れ。

- Abe, K., 2002: Preformed Cd and PO<sub>4</sub> and the relationship between the two elements in the northwestern Pacific and the Okhotsk Sea. *Mar. Chem.*, **279**, 27-36.
- Abe, K., 2002: Relationship between Cd and PO<sub>4</sub> in the subtropical sea near the Ryukyu Islands. *J. Oceanogra.*, **58**, 577-588.
- Abe, K., 2002: Preformed Cd and PO<sub>4</sub> in the northeastern Pacific and the Okhotsk Sea. *West. Pac. Geophys. Meet.*, pp.37.
- Abe, K., 2002: Preformed Cd and PO<sub>4</sub> in the equatorial Pacific deep water. Abstract of the 2002 Japan Earth and Planetary Science Joint Meeting, Tokyo,(cd-rom).
- Abe, K., 2003: Dissolved copper in the Yellow Sea and the East China Sea-Cu as a tracer of the Changjiang discharge -. *Deep Sea Res. II*, **50(2)**, 327-337.
- Abe, K., 2003: Kink in the Cd-PO<sub>4</sub> plot near the Okinawa Island observed in the summer of 2001. *Bull. Fish. Res. Agen.*, **6**, 15-17.
- Abe O., Shibuno T., Takada Y., and Hashimoto K., 2002: Preliminary study to prevent incidental capture of sea turtles by the pound net. *SEASTAR2000*, pp.29
- Abe S., Iguchi K., Ito S., Uchida Y., Ohnishi H., and Ohmori K., 2002: Habitat use of the grazing goby (*Sicyopterus japonicus*). *J. Freshwater Ecol.*, **18**, 161-167.
- 阿部信一郎，南雲 保，田中次郎，2002: ロジスティックモデルによる河川付着藻類群落の量的動態の解析。陸水学雑誌，**63**，209-213。
- Adachi M., Matsubara T., Okamoto R., Nishijima T., Itakura S., Yamaguchi M., 2002: Inhibition of cyst formation in the toxic dinoflagellate *Alexandrium* (Dinophyceae) by bacteria from Hiroshima Bay Japan. *Aquat. Microb. Ecol.*, **26**, 223-233.
- 阿保勝之，坂見知子，高柳和史，2002: アサリ増殖場造成地における水質浄化機能の定量的評価。海岸工学論文集，**46(2)**，1211-1215。
- 赤羽敬子，岸 道郎，向井 宏，飯泉 仁，2003: 陸域からの栄養塩負荷量に対する北海道厚岸湖の生態系の応答。沿岸海洋研究，**40(2)**，171-179。
- Akamatsu T., Okumura T., Novarini N., and Yan H. Y., 2002 : Empirical refinements applicable to the recording of fish sounds in small tanks. *J. Acoust. Soc. Am.*, **112**, 3073-3082.
- Akamatsu T., Wang D., Wang K., Wei Z., Q.Zhao Q., and Naito Y., 2002: Diving behaviour of freshwater finless porpoises (*Neophocaena phocaenoides*) in an oxbow of the Yangtze River, China. *ICES J. Mar. Sci.*, **59**, 438-443.
- Akamatsu T., Wang D., and Wang K., Wei Z., and Naito Y., 2002: A passive acoustical survey method of finless porpoises. *Fish. Sci.*, **68(Suppl. )**, 294-297.
- 赤松友成，2002: 講座（海洋計測）：(1) クジラとイルカを声で見る。海洋音響学会誌，29， 61-66。
- 赤松友成，2003: イルカは音で水中を探る。科学，**73(1)**，18-22。
- 赤嶺達郎，平松一彦，2003: 資源管理の研究。日水誌，**69**特別号，51-58。
- Akeda S., Nakamura Y., and Takaki N., 2002 : Designing the environmentally conscious projects for coastal development. *PACON-2002*, **S7B**, pp.99.
- Aketa S., Takaki N., and Iseki K., 2002 : Possibilities of creating new fishing grounds using deep seawaters. *PACON-2002*, pp.206.
- 明田定満，桑原久実，中村義治，奥出 壮，寺澤知彦，2002: エゾアワビの個体群動態モデルの開発。海岸工学論文集，**49**，1171-1175。

- 秋山秀樹, 2003: 南西海区における早期海況変動予測システム . 黒潮の資源海洋研究 , 4, 69-71.
- Amano K., Ookubo N., Ikuta K., Kitamura S., and Okuzawa K., 2002: Ontogenetic origin of salmon GnRH neurons in the ventral telencephalon and the preoptic area in masu salmon. *Gen. Comp. Endocrinol.*, **127** (3), 256-262.
- Anai M., Park H.C., Saeki K., Akeda S., Yamamoto K., Kondo T., and Takaki N., 2002 : Analytical study on the fishermen's working assessment. *Techno-Ocean2002*, (cd-rom)
- 穴井美緒, 朴 賢哲, 佐伯公康, 明田定満, 近藤健雄, 高木伸雄, 2003: 漁港における漁業就労環境の基礎的研究 . 日本沿岸域学会論文集 , **15**, 113-120..
- Ando W., Nakayama A., Kuwahara H., and Kawamata S., 2002: The fishing port construction harmonized with the ecosystem of making seaweed forests. *Techno-Ocean 2002*, s-v-7 (cd-rom)
- Ayaz Z., Vassalos D., Spyrou K. J., and Matsuda A., 2002 : Towards a unified 6 DOF manoeuvring model in random seas. *Proc. 6th Internl. Ship Stability Workshop*, 3.2.1-3.2.9.
- 東 幹夫, 斎藤憲治, 片野 修, 2002: 魚類の調査方法-モンドリによる淡水魚の採捕 . 魚類学雑誌 , 49, 158-160.
- Azuma T., Noda S., Yada T., Ototake M., Nagoya H., Moriyama S., Yamada H., Nakanishi T., and Iwata M., 2002: Profiles in growth, smoltification, immune function and swimming performance of 1-year-old masu salmon (*Oncorhynchus masou masou*) reared in water flow *Fish. Sci.*, **68**(6), 1282-1294.
- Azuma T., Ohta H., Oda S., Muto K., Yada T., and Unuma T., 2003: Changes in fertility of rainbow trout eggs retained in coelom. *Fish. Sci.*, **69**(1), 131-136.
- Brodeur R. D., Sugisaki H., and Hunt G. L., 2002 : Increases in jellyfish biomass in the Bering Sea: implications for the ecosystem. *Mar. Ecol. Prog. Ser.*, **233**, 89-103.
- Brownell R. L. Jr., Clapham P., Miyashita T. and Kasuya T., 2001: Conservation status of North Pacific right whales. *J. Cetacean Res. Manag. Special Issue*, **2**, 269-286.
- Dijkstra J.M., Yoshiura Y., Kiryuu I., Aoyagi K., and Koellner B., 2003: The promoter of the classical MHC class I locus in rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.*, **14**, 177-185.
- Fernandez-Leborans G., Hanamura Y., and Nagasaki K., 2002: A new suctorian, *Flectacinetes isopodoensis* (Protozoa: Ciliophora) epibiont on marine isopods from Hokkaido (northern Japan). *Acta Protozoo.*, **141**, 79-84.
- Fischer U., Utke, K., Ototake M., Dijkstra J.M., and Koellner B., 2003: Adaptive cell-mediated cytotoxicity against allogeneic targets by CD8-positive lymphocytes of rainbow trout (*Oncorhynchus mykiss*). *Develop. Comp. Immunol.*, **26**, 131-139.
- 藤井一則, 角埜 彰, 原 彰彦, 2002: 時間分解蛍光免疫測定法によるマコガレイ *Pleuronectes yokohamae* コリオジエンの定量法開発 . 環境毒性学会誌 , **5**, 33-41.
- Fujiwara A., Kiryuu I., Dijkstra J.M., Yoshiura Y., Nishida-Umebara C., 2003: Chromosome mapping of MHC class I in rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.*, **14**, 171-175.
- Fujiyoshi, E., 2002: Cryopreservation of conchocelis of *Porphyra yezoensis* form. *narawaensis* without DMSO. *1st Cong. Internat. Soc. Appl. Algol.*, 2002, pp.108.
- \*Fukuda M., Sako H., Shigeta T.,and Shibata.,2002 : Effects of body size and temperature on the relationship between growth and ration of young Japanese flounder, based laboratory experiments. *Bull. Fish. Res. Agen.*, **4**, 11-18.
- 福田光男, 坪田幸雄, 森 信幸, 丸山修治, 吉田 徹, 久野能孝, 山下俊彦, 2002: 底質・流速からみた沿岸構造物周辺の底生生物分布の経年変化 . 海岸工学論文集 , **49**, 1221-1225.
- 福田 穩, 岡村 愛, 西山 勝, 川上秀昌, 釜石 隆, 2002: 養殖イサキに発生した細胞内寄生細菌による肉芽腫症 . 魚病研究 , **37**(3), 119-124.
- Fukuoka K., Hanamura Y., and Murano M.,2002 : Supplementary information on the taxonomy and distribution of six species of Anisomysis (Crustacea: Mysidacea: Mysidae). *Plankton Biol. Ecol.*, **49**, 38-43.
- 福富則夫, 中村智幸, 土居隆秀, 武田維倫, 尾田紀夫, 2002: 栃木県那珂川水系における ミツバヤツメの捕獲の記録 . 魚類学雑誌 , **49**(1), 53-58.

- \*Fukuwaka M. and Suzuki T., 2002: Early sea mortality of mark-recaptured juvenile chum salmon in open coastal waters. *J. Fish Biol.*, **60**(1), 3-12.
- Fukuwaka M., and Suzuki T., 2002: Early sea mortality of mark-recaptured juvenile chum salmon in open coastal waters. *J. Fish Biol.*, **60**(1), 5-12.
- Furuita H., Tanaka H., Yamamoto T., Suzuki N., and Takeuchi T., 2002: Effects of high levels of n-3 HUFA in broodstock diet on egg quality and egg fatty acid composition of Japanese flounder. *Aquaculture*, **210**, 323-333.
- Gen K., Okuzawa K., Tanaka H., and Kagawa H., 2002: Sexual dimorphism of gonadotropin Ib gene expression in red seabream, *Pagrus major*. *Fish. Sci.*, **68**(Suppl. ), 671-674.
- 玄 浩一郎 , 2002: 魚類の性成熟に関する視床下部 - 脳下垂体系に関する研究 . 日水誌 , **68**(4),497-499.
- Goto T., 2002: Changes in the distribution and abundance of *Todarodes pacificus* (Cephalopoda, Ommastrephidae) paralarvae in the southwest Sea of Japan with changing stock levels. *Fish. Sci.*, **68**(Suppl. ), 198-201.
- Haga Y., Suzuki T., and Takeuchi T., 2002: Retinoic acid isomers produce malformations in postembryonic development of the Japanese flounder, *Paralichthys olivaceus*. *Zool. Sci.*, **19**(10), 1105-1112.
- Hakoyama H., 2003: The ideal free distribution when the resource is variable. *Behav. Ecol.*, **14**, 47-53.
- 浜口昌巳 , 佐々木美穂 , 薄 浩則 , 2002: 日本国内におけるアサリ *Ruditapes philippinarum* の *Perkinsus* 原虫の 感染状況 . 日本ベントス学会誌 , **57**, 168-178.
- 浜口昌巳 , 佐々木美穂 , 2002: 発生初期のアワビ類を判別する技術の開発 . 月刊海洋 , **34**, 517-521.
- Hamano k. and Awaji M., 2002 : cDNA cloning and expression analysis of insulin-related peptide gene and prohormone convertase gene in the Pacific oyster, *Crassostrea gigas*. *Fish. Sci.*, **68**(Suppl. ), 769-771.
- 原 素之, 関野正志, 2002:アワビ栽培漁業におけるマイクロサテライトDNA分析の応用. 月刊海洋 , **34**(7), 512-516.
- Hara M. and Sekino M., 2003 : Efficient detection of parentage in a cultured Japanese flounder *Paralichthys olivaceus* using microsatellite DNA markers. *Aquaculture*, **217**, 107-114.
- 長谷川勝男 , 井原雄一 , 2002: 小型高速機関の低温始動性における燃料油性状の影響 . 日本マリンエンジニアリング 学会誌 , **37**, 24-30.
- Hashimoto H., Umeda N., and Matsuda A., 2002 : Enhanced approach for broaching prediction with higher order terms taken into account. *Proc. 6th Internl. Ship Stability Workshop*, 3.3.1-3.3.8.
- 林原 毅 , 2002: 造礁性イシサンゴ類の配偶子放出の周期性と同調機構 . 海洋と生物 , **24**(6), 491-498.
- Hewitt R. P., Watkins J. L., Naganobu M., Tshernyshkov P., Brierley A. S., Demer D., Kasatkina S., Takao Y., Goss C., Malyshko A., Brandon M. A., Kawaguchi S., Siegel V., Trathan P. N., Emery J. H., Everson I., and Miller D. G. M., 2002: Setting a precautionary catch limit for antarctic Krill. *Oceanography*, **15**(3), 26-33.
- Higano J. and Pichitkul P., 2002: A nutrient budget in some polyculture systems of giant tiger prawn and green mussel in Thailand. *Fish. Sci.*, **68**(Suppl. ), 839-842.
- 平松一彦 , 岡村 寛 , 2003: 小標本における標準誤差と信頼区間の推定について . 水研センター研報 , **6**, 1-6.
- Hiroe Y., Yasuda I., Komatsu K., Kawasaki K., Joyce T. M., and Bahr F., 2002 : Transport of North Pacific intermediate water in the Kuroshio-Oyashio interfrontal zone. *Deep-Sea Res.* , **49**: 5353-5364.
- Hiyama,Y., Yoda,M., and Oshima, S.,2002: Stock size fluctuations in chub mackerel (*Scomber japonicus*) in the East China Sea and the Japan/East Sea. *Fish. Oceanogr.*, **11**, 347-353.
- Honda, H., Ishida M., Mitani T., Uehara S., Hirota Y., Sakaji H., Nashida K., and Morimoto H., 2002 : Recent trends in age structures and maturity conditions of the Japanese sardine, *Sardinops melanostictus*, spawning in Tosa Bay. *Fish. Sci.*, **68**(Suppl. ), 210-213.
- Hoshika A., Tanimoto T., Mishima Y., Iseki K., and Okamura K., 2002: Variation of turbidity and particle transport in the bottom layer of the East China Sea. *Deep-Sea Res.*, **50**, 443-455.
- Hu F., Oozeki Y., Tokai T., and Matsuda K., 2002: Development of a new midwater sampling trawl. *Fish. Sci.*, **68**(Suppl. ), 1899-1900.

- Ichimi K., Suzuki T., and Itoh A., 2002: Variety of PSP toxin profiles in various culture strains of *Alexandrium tamarense* and change of toxin profile in natural A-tamarense population. *J. Exp. Mar. Biol. Ecol.*, **273**(1), 51-60.
- Ichii T., Mahapatra K., Watanabe T., Yatus A., Inagake D., and Okada Y., 2002: Occurrence of jumbo flying squid *Dosidicus gigas* aggregations associated with the countercurrent ridge off the Costa Rica Dome during 1997 El Nino and 1999 La Nin?. *Mar. Ecol. Prog. Ser.*, **231**, 151-166.
- Ichikawa Y., Takatsuki Y., Mizuno K., Shikama N., and Takeuchi K., 2002: Estimation of drifting velocity and error at parking depth for the Argo float. *ARGO Technical Report FY2001*, JAMSTEC, pp.68-77.
- 市川忠史, 加藤 聰, 濑川恭平, 2003: プランクトンセンサーで何が分かるか - 水産海洋学の視点から - . 日本プランクトン学会報, **50**(1), 29-35.
- Iguchi K., Itoh F., Ogawa K., Matsubara N., Yodo T., and Yamasaki T., 2002: Reduction of transport stress of ayu by obligated schooling. *Fish. Sci.*, **68**(4), 849-853.
- Iguchi K., Ogawa K., Nagae M., and Ito F., 2002: The influence of rearing density on stress responded disease susceptibility of ayu (*Plecoglossus altivelis*). *Aquaculture*, **220**, 515-523.
- Iguchi K. and Abe S., 2002: Territorial defense of an excess food supply by an algal grazing fish ayu. *Ecol. Res.*, **17**, 373-380.
- Iguchi K. and Nishida M., 2002: Genetic biogeography among insular populations of ayu. *Fish. Sci.*, **68**(Suppl.), 345-348.
- 飯泉 仁, 鈴木健吾, 2002: 厚岸湖生態系における安定同位体比の分布. 月刊海洋, **13**(6), 412-416.
- 池田久美子, 南 卓志, 山田 久, 小山次朗, 2002: 日本海底層の食物網における有機スズ化合物の生物濃縮. 環境化学, **12**(1), 105-114.
- Ikeda K., Kidokoro H., Arai N., and Sakamoto W., 2003: Strontium:calcium ratios in statoliths of Japanese common squid *Todarodes pacificus* (Cephalopoda: Ommastrephidae) as indicators of migratory behavior. *Mar. Ecol. Prog. Ser.*, **251**, 169-179.
- Ikeda Y., Yatsu A., Arai N., and Sakamoto W., 2002: Concentration of statolith trace elements in the jumbo flying squid during El Nino and non-El Nino years in the eastern Pacific. *J. Mar. Biol. Ass. U.K.*, **82**, 863-866.
- Imamura S., Ojima N., and Yamashita M., 2002: Molecular cloning and cold-inducible gene expression of the cell division cycle gene CDC48 in zebrafish cells. *Fish. Sci.*, **68**(Suppl.), 1291-1292.
- 稻掛伝三・植原量行 (2003): まぐろ類の資源変動と大気／海洋変動. 月刊海洋, **35**, 180-195.
- Inagawa H., Honda A., Kawauchi C., Nishizawa T., and Yoshiura Y., 2002: Cloning and characterization of the homolog of mammalian lipopolysaccharide-binding protein and bactericidal permeability-increasing protein in rainbow trout *Oncorhynchus mykiss*. *J. Immunol.*, **168**, 5638-5644.
- Inoue A., Miyazaki M., Izawa K., Ando K., Takatsuki Y., and Mizuno K., 2002: Stability of water temperature in the conductivity and temperature calibration system and results of calibration experiments. *ARGO Technical Report FY2001*, JAMSTEC, pp. 9-17.
- Iseki K., Terawaki T., Yoshikawa K., Yoshida G., Sakanishi Y., Yoshimura T., Kawamata S., Kuwahara H., and Noda M., 2002: Evaluation of global warming impacts to seagrass and seaweed beds; An introduction of "Global Warming Program" of the Fisheries Research Agency. *Techno-Ocean 2002*, s-v-1(cd-rom).
- Iseki K., Okamura K., and Kiyomoto Y., 2002: Seasonality and composition of downward particulate fluxes at the continental shelf and Okinawa Trough in the East China Sea. *Deep-Sea Res.*, **50**, 457-473.
- 井関和夫, 2002: 散布深層水の挙動把握と海域肥沃化に関する研究: 漂流ブイと自動昇降型C TD・クロロフィル計の有効性について. 海洋深層水研究, **3**(2), 83-90.
- 井関和夫, 2002: 懸濁粒子および沈降粒子の採取. 海洋調査, **71**, 6-12.
- Ishida Y., Azumaya T., Fukuwaka M., and Davis N., 2002: Interannual variability in stock abundance and body size of Pacific salmon in the central Bering Sea. *Prog. Oceanogr.*, **55**(1-2), 223-234.
- Ishihara K., Komatsu W., Saitoh H., and Shinohara K., 2002: Comparison of the effects of dietary-linolenic,

- stearidonic and eicosapentaenoic acids on production of inflammatory mediators in mice. *Lipids*, **37(5)**, 481-486.
- Ishii K., Miyahana Y., and Sawada K., 2002: Transducer rotation system for target strength measurement of cephalopod. *Proceedings of International Symposium on Advanced Techniques of Sampling Gear and Acoustical Surveys for Estimation of Fish Abundance and Behavior (ACOUSTGEARE 2000)*, pp.30-34.
- Ito L. S., Yamashita M., and Strussman C., 2002: Dynamics of heat-induced germ cell loss in pejerrey, *Odontesthes bonariensis*. *Fish. Sci.*, **68(Suppl. )**, 1313-1314.
- Iwasaka N., Fujita T., and Mizuno K., 2002: Research and development of the method for the deployment of the Argo float. *JAMSTECR*, **46**, 81-94.
- 岩崎俊秀, 2002: 6年もかかりました。電波受験界, **50**, 12-17.
- Iwasaka N., Suga T., Takeuchi K., Mizuno K., Takatsuki Y., Ando K., Kobayashi T., Oka E., Ichikawa Y., Miyazaki M., Matsuura H., Izawa K., Yang C.-S., Shikama N., and Aoshima M., 2003: Pre-Japan-ARGO: Experimental observation of upper and middle layers south of the Kuroshio Extension region by using profiling floats. *J. Oceanogr.*, **59**, 119-127.
- Izawa K., Mizuno K., Miyazaki M., Inoue A., Ando K., Takatsuki Y., Kobayashi T., and Takeuchi K., 2002: On the weight adjustment of profiling floats. *ARGO Technical Report FY2001*, JAMSTEC, pp. 18-35.
- Jinnai K., Watanabe Y., Sugiyama T., Ootsuki T., Ooide A., Azuma T., Iwata M., Okino T., and Yayagisawa T., 2002: Hormonal existence of thyroid axis in sockeye salmon (*Oncorhynchus nerka*) during down-river migration. *Fish. Sci.*, **68(Suppl. )**, 961-962.
- Kamaishi T. and Yoshinaga T., 2002: Detection of *Haplosporidium nelsoni* in Pacific oyster *Crassostrea gigas* in Japan. *Fish Pathol.*, **37**, 193-195.
- Kamisaka Y., Kaji T., Masuma S., Tezuka N., and Kurokawa T., Suzuki T., Totland G. K., Ronnestad I., Tagawa M., and Tanaka M., 2002: Ontogeny of cholecystokinin-immunoreactive cells in the digestive tract of blue fin tuna, *Thunnus thynnus*, larvae. *Sarsia*, **87**, 258-262.
- Katano O., Yamamoto S., and Nakamura T., 2002: Predation of Japanese dace, *Tribolodon hakonensis*, by largemouth bass, *Micropterus salmoides*, in experimental aquaria. *Ichthyol. Res.*, **49**, 392-396.
- Katano O. and Aonuma Y., 2002: An experimental study of the effects of smallmouth bass on the behavior, growth and survival of prey fish. *Fish. Sci.*, **68(5)**, 803-814.
- Katano O., Yamamoto S., and Nakamura T., 2002: Predation of Japanese dace, *Tribolodon hakonensis* by large mouth bass, *Micropterus salmoides*, in experimental aquaria. *Ichthyol. Res.*, **49**, 392-396.
- Kato H. and Kasuya T., 2002: Some analyses on the modern whaling catch history of the western North Pacific stock of gray whales (*Eschrichtius robustus*), with special reference to the Ulsan whaling ground1. *J. Cetacean Res. Manag.*, **4**, 277-282.
- Kawai H., Yatsu A., Watanabe C., Mitani T., Katsukawa T., and Matsuda H., 2002: Recovery policy of chub mackerel stock using recruitment-per-spawning. *Fish. Sci.*, **68(5)**, 963-971.
- 川井唯史, 金田友紀, 新井章吾, 桑原久実, 2002: 磯焼け地帯におけるウニ侵入防止フェンスによるホソメコンブ群落の造成とキタムラサキウニ生殖巣の発達. *水産工学*, **39(1)**, 15-20.
- 川西 澄, 水野博史, 松山幸彦, 長井 敏, 小谷祐一, 2002: 広島湾北部域における底層懸濁粒子からの音響散乱特性 - 超音波ドップラー分布計による底層濁度と体積濃度の測定可能性について. *海の研究*, **11(2)**, 285-293.
- 川西 澄, 松山幸彦, 2002: 内湾域に存在する海中懸濁粒子からの音響散乱 *海岸工学論 文集*, **49**, 991-995.
- Kawasaki H., Sano M., and Shibuno T., 2003: The relationship between habitat physical complexity and recruitment of the coral reef damselfish, *Pomacentrus amboinensis*: and experimental study using smallscale artificial reefs. *Ichthyol. Res.*, 2003, 73-77.
- Kawashima T., Yoshimura Y., and Suzuki S., 2002: Application of Rankine source method for improving hull form of fishing vessels. *Fish. Sci.*, **68(Suppl. )**, 1807-1810.
- Kim D., Sato Y., Miyazaki Y., Oda T., Muramatsu T., Matsuyama Y., and Honjo T., 2002: Comparison of hemolytic activities among strains of *Heterocapsa circularisquama* isolated in various localities in

- Japan. *Biosci. Biotechnol. Biochem.*, **66**(2), 453-457.
- Kim D., Oda T., Muramatsu T., Kim D., Matsuyama Y., and Honjo T., 2002: Possible factors responsible for the toxicity of *Cochlodinium polykrikoides*, a red tide phytoplankton. *Comp. Biochem. Physiol. C*, **132**(4), 415-423.
- Kim S.-K., Takeuchi T., Akimoto A., Furuita H., and Murata Y., 2002: The effect of a fishmeal-based diet supplemented with taurine on juvenile Japanese flounder *Paralichthys olivaceus*. *Fish. Sci.*, **68**(Suppl. ), 989-990.
- 木村 量, 梨田一也, 大関芳沖, 本多 仁, 2002: ゴマサバ *Scomber australasicus*に適した耳石による年齢査定法 . 水産海洋研究 , **66**(4), 247-251.
- 木村 量, 梨田一也, 大関芳沖, 本多 仁, 2002: ゴマサバ耳石の年齢を読む試み . 黒潮の資源海洋研究 , 3, 93-96.
- Kitagawa T., Nakata H., Kimura S., Sugimoto T., and Yamada H., 2002: Differences in vertical distribution and movement of Pacific bluefin tuna (*Thunnus thynnus orientalis*) among areas: the East China Sea, the Sea of Japan and the western North Pacific. *Mar. Freshwat. Res.*, **53**, 245-252.
- Kitagawa T., Nakata H., Kimura S., Sugimoto T., and Yamada H., 2002: Analysis of tunas behavior using acoustic, archival and pop-up tags: a review. *Otsuchi Mar. Sci.*, **27**, 1-6.
- Kitagawa T., Nakata S., Kimura H., and Yamada H., 2002: Diving behavior of immature Pacific bluefin tuna (*Thunnus thynnus orientalis*) recorded by an archival tag. *Fish. Sci.*, **68**(Suppl. ), 427-428.
- 清田雅史, 2002: 延縄漁業における海鳥類の偶發的捕獲: 問題の特性と回避の方法 . 山階鳥研報 , 34, 1-19.
- Kobayashi T., Hosoda S., Suga T., Mizuno K., Shikama N., and Takeuchi K., 2002: Ocean conditions around southeast off Japan estimates by the Argo data with optimum interpolation method. *JAMSTECR*, **46**, 1-9.
- Kobayashi T., Nakajima H., Suga T., Mizuno K., Shikama N., and Takeuchi K., 2002: High quality climatological dataset for the Indian Ocean (Indian HydroBase). *JAMSTECR*, **46**, 11-27.
- Kobayashi T., Ichikawa Y., Takatsuki Y., Suga T., Iwasaka N., Ando K., Mizuno K., Shikama N., and Takeuchi K., 2002: Quality control of Argo data based on high quality climatological dataset (HydroBase) I. *ARGO Technical Report FY2001*, JAMSTEC, pp.38-48.
- Kobayashi T., Ichikawa Y., Takatsuki Y., Suga T., Iwasaka N., Ando K., Mizuno K., Shikama N., and Takeuchi K., 2002: Correction method for Argo data based on HydroBase I-Introduction of potential conductivity -. *ARGO Technical Report FY2001*, JAMSTEC, pp.49-56.
- Kobayashi T., Ichikawa Y., Takatsuki Y., Suga T., Mizuno K., Iwasaka N., Shikama N. and Takeuchi K., 2002: Study of density range with seasonal variations in water-mass structure. *ARGO Technical Report FY2001*, JAMSTEC, pp.57-67.
- Kogure Y., 2002: First records of the brooding astropectinid starfish *Trophodiscus almus* (Echinodermata: Asteroidea) from Japan, with reference to its zoogeographical significance. *Species Diversity*, **7**(2), 187-203.
- 小松幸生, 川崎 清, 2002: 黒潮下流域遠州灘沖合特有の中層水構造 . 水研センター研報 , 5, 1-22.
- 小松幸生, 友定 彰, 川崎 清, 2002: 伊豆小笠原海嶺域における定期船航走水温の時空間変動特性 . 年々変動 . 水研センター研報 , 5, 23-37.
- 小松幸生, 友定 彰, 川崎 清, 2002: 伊豆小笠原海嶺域における定期船航走水温の時空間変動特性 . 年内変動 . 水研センター研報 , 5, 38-54.
- 小松幸生, 川崎 清, 2003: 漂流ブイによる東シナ海表面吹送流の変動特性 . 水研センター研報 , 7, 14-23.
- 小西光一, 2002: 幼生研究のための小テクニック集 (3) 文献の収集と整理. *Cancer*, **11**, 41-47.
- Kosuge, T., and Maung, M. T., 2002: Status of mud crab fishery in Ayeyarwady mangroves, Myanmar 2000-2001. Proceedings of Seminar on Mangrove Reforestation through Community Participation, pp.1-10.
- Kubota H., Tsuchiyama A., Oozeki Y., Kimura R., Kado R., Ogawa H., Hu F., and Matsuda K., 2002: Diel vertical distribution of larvae and juveniles of pelagic and mesopelagic fishes by a newly developed

- frame trawl in the Kuroshio-Oyashio Transition Area in spring. *Fish. Sci.*, **68(Suppl. )**, 1897-1898.
- Kurata K., Minami H., and Kikuchi E., 2001: Stable isotope analysis of food sources for salt marsh snails. *Mar. Ecol. Prog. Ser.*, **223**, 167-177.
- Kurihara, T., 2002: Spatial and temporal fluctuation in the density of the intertidal limpet, *Patelloidea striata* Quoy et Gaimard, on subtropical cobbled shores. *J. Molluscan Stud.*, **68**, 83-90
- 栗田 潤, 飯田悦左, 中島員洋, 井上 潔, 2002: ヒラメVHSウイルスに対する各種市販消毒剤の殺ウイルス効果 . 魚病研究 , **37(4)**, 175-181.
- Kurogi H., Mochioka N., Takagi Y., and Tabeta O., 2002: First offshore record of *Conger myriaster leptocephali* in the East China. *Fish. Sci.*, **68(5)**, 1155-1157.
- Kurokawa T., Suzuki T., Ohta H., Kagawa H., Tanaka H., and Unuma T., 2002: Expression of pancreatic enzyme genes during the early larval stage of Japanese eel *Anguilla japonica*. *Fish. Sci.*, **68(4)**, 736-744.
- Kurokawa T. and Suzuki T., 2002: Development of neuropeptide Y-related peptides in the digestive organs during the larval stage of Japanese flounder, *Paralichthys olivaceus*. *Gen. Comp. Endocrinol.*, **126**, 76-79.
- Kuwahara H., 2002: The fishing port construction harmonized with the ecosystem of making seaweed forests. *Techno-Ocean 2002*, s-v-7(cd-rom)
- Kuwahara H., Kaneda T., and Kawai T., 2002: Engineering methods for kelp forest development on the barren ground along the southwest coast of Hokkaido, Japan. *Fish. Sci.*, **68(Suppl. )**, 1751-1754.
- Kuwahara H., Akeda S., and Sakou A., 2002: Assessment method for developing sea urchin fishing grounds and sea algae bed using the population dynamics model. *Techno-Ocean 2002*, s-v-6 (cd-rom).
- Kuwahara H., Akeda S., and Sakou A., 2002: The development of technique for evalaating sea-uchi fishing grounds using the polalation dynamics model. *3rd CST panel of UJNR*, pp.113-133.
- 桑原久実, 川井唯史, 金田友紀, 2002: 磯焼け海域の藻場造成礁におけるホソメコンブ群落を維持するために必要な流動条件 . 水産工学 , **39**, 47-53.
- 桑原久実, 明田定満, 酒向章哲, 2002: 個体群動態モデルを用いたウニ漁場評価手法の開発 . 海岸工学論文集 , **49**, 1246-1250.
- Leedom, T.A., Uchida K., Yada T., Richman N. H., and Byatt J.C., 2002: Recombinant bovine growth hormone treatment of tilapia: growth response, metabolic clearance, receptor binding and immunoglobulin production. *Aquaculture*, **207(3-4)**, 359-380.
- Liang X. F., Ogata H., and Oku H., 2002: Effect of dietary fatty acids on lipoprotein lipase gene expression in the liver and visceral adipose tissue of fed and starved red sea bream *Pagrus major*. *Comp. Biochem. Physiol.A*, **132**, 913-919.
- Mackenzie L., Holland P., McNabb P., Beuzenberg V., Selwood A., and Suzuki T., 2002: Complex toxin profiles in phytoplankton and Greenshell mussels (*Perna canaliculus*), revealed by LC-MS/MS analysis. *Toxicon*, **40(9)**, 1321-1330.
- Mackenzie L., Suzuki T., and Adamson J., 2001: Elimination and differential transformation of yessotoxin by the greenshell mussel *Perna canaliculus* and the blue mussel *Mytilus galloprovincialis*. *Proc. 9th Intl. Conf. Harmful Algal Blooms*, 371-374.
- 馬 寧, 芳村康男, 木村暢夫, 鈴木四郎, 山越康行, 2002: バルジによる改造漁船船型の波浪中運動の基本特性 . 日本造船学会論文集 , **192**, 219-225.
- 丸山修治, 福田光男, 森 信幸, 坪田幸雄, 2002: 溶存酸素収支に基づく蓄養施設一体型構造物の蓄養可能量 . 海洋論文集 , **18**, 527-532.
- 正岡哲治, 小林敬典, 2002: 18SrRNA及び28SrRNA領域を用いたアコヤガイ属における系統関係 . DNA多型 , **10**, 100-104.
- 升間主計, 手塚信弘, 尾花博幸, 鈴木伸明, 野原健司, 張 成年, 2003: ミトコンドリアDNA分析から推定した養成クロマグロの産卵生態 . 水研センター研報 , **6**, 9-14.
- 松浦 勉, 2002: 太平洋南区における沖合底びき網の経営動向の分析 . 北日本漁業 , **30**, 65 -77.
- 松浦 勉, 2002: 定置漁村と観光定置の成立条件の関係について . 水研センター研報 , **5**, 55-67.

- 松浦 勉, 2003: 鳥取県浜田市における沖合底びき網漁業の経営と就労動向の分析 . 北日本漁業 , **31**, 103-121.
- Miyahana Y., Sawada K., Takao Y., Kuratsu K., and Satoh T., 2002: Quantitative echo sounding system for underwater robots. *Proceedings of International Symposium on Advanced Techniques of Sampling Gear and Acoustical Surveys for Estimation of Fish Abundance and Behavior (ACOUSTGEARE 2000)*, pp.55-60.
- Masuya T., 2002: An application of unconventional hull form to fishing vessels. *Fish. Sci.*, **68(Suppl. )**, 1811-1814.
- 松田秋彦, 橋本博公, 梅田直哉, 2002: バウダイビングによる転覆現象ー第一報 バウダイビング発生条件ー . 日本造船学会論文集 , **192**, 201-207.
- Matsuda A., Umeda N., and Hashimoto H., 2002: Comparison of North European and Japanese Purse Seiners by capsizing model experiments. *J. Kansai Soc. Naval Architects Japan*, **238**, 121-129.
- Matsuda A., Umeda N., Hashimoto H., and Urano S., 2002: Comparison of North European and Japanese Purse Seiners by capsizing model experiments. *Proc. 1st. Asia Pacific Workshop Mar. Hydrodynamics*, pp.50-54.
- Matsuda A., Umeda N., and Suzuki S., 2002: Capsizing experiments of purse seiners in following and quartering seas and anti-broaching steering system. *Fish. Sci.*, **68(Suppl. )**, 1831-1834.
- Matsuoka M., 2002: Review:Comparison of meristic variations and bone abnormalities between wild and laboratory-reared red sea bream. *Jpn. Agri. Res. Quart.*, **37(1)**, 21-30.
- 松岡正信, 2002: コモンフグ天然魚にみられた鼻孔隔皮異常について . 水産増殖 **50**, 233 -234.
- 松岡正信, 2002: 人工採苗クロダイにおける鼻孔隔皮と嗅板の形成過程 . 水産増殖 , **50(4)**, 401-406.
- 松岡正信, 2002: 鼻孔隔皮欠損症の発生率が比較的低い飼育例におけるマダイの鼻孔隔皮 発達過程 . 水産増殖 , **51(1)**, 121-122.
- Matsumoto T., Nakamura A., Mori K., and Kayano T., 2003: Molecular characterization of a cDNA encoding putative vitellogenin from the Pacific oyster *Crassostrea gigas*. *Zool. Sci.*, **20(1)**, 37-42.
- Matsushita Y., Glass C.W., Morris G. D., Feehan T. L., and Sarno B., 2002: Taking fishermen's knowledge into conservation harvesting technology research and development-a case study in New England, USA-. *Fish. Sci.*, **68(Suppl. )**, 376-379.
- Matsuyama Y., Uchida T., Honjo T., and Shumway S.E., 2002: Impacts of the harmful dinoflagellate, *Heterocapsa circularis quama*, on shellfish aquaculture in Japan. *J. Shellfish Res.*, **20(3)**, 1269-1272.
- 南川真吾, 2003: 動物の潜水速度はどうやって決まるのか. 科学, **173**, 54-55.
- Miyake H., Lindsay D. J., Hunt J. C., and Hamatsu T., 2002: Scyphomedusa *Aurelia limbata* (Brandt, 1838) found in deep waters off Kushiro, Hokkaido, northern Japan. *J. Plankton Biol. Ecol.*, **49(1)**, 44-46.
- Miyamoto H., Hamaguchi M., and Okoshi K., 2002: Analysis of genes expressed in the mantle of oyster *J. Crassostrea gigas*. *Fish. Res.*, **68(3)**, 651-658.
- 宮本政秀, 吉田雄一, 河邊 博, 松山幸彦, 高山晴義, 2002: 1995年に羊角湾で発生した渦鞭毛藻*J. Gyrodinium* sp. の赤潮: 発生期の環境特性と養殖魚への影響 . 日水誌 , **68(2)**, 157-163.
- Mizobata K., Saitho S., Shiromoto A., Miyamura T., Shiga N., Imai K., Toratani M., Kajiwara Y., and Sasaoka K., 2002: Bering Sea cyclic and anticyclonic eddies observed during summer 2000 and 2001. *Prog. Oceanogr.*, **55(1-2)**, 65-76.
- Mizuno K., 2003: On the termination of Argo float. *Umi no Kenkyu*, **12**, 1-21.
- Mochida K., Matsubara T., Andoh T., Ura K., Adachi S., and Yamauchi K., 2002: A novel seminal plasma glycoprotein of a teleost, the Nile tilapia (*Oreochromis niloticus*), contains a partial von Willebrand factor type D domain and a zona pellucida like domain. *Mol. Reprod. Develop.*, **62**, 57-68.
- Mochida K., Matsubara T., Kudo H., Andoh T., Ueda H., Adachi S., and Yamauchi K., 2002: Molecular cloning and immunohistochemical localization of ubiquitin C-terminal hydrolase expressed in testis of a teleost, the Nile tilapia, *Oreochromis niloticus*. *J. Exp. Zool.* **293**, 368-383.
- Mochida K. and Matsubara T., 2002: Molecular cloning of an elongation factor 1a and its mRNA localization

- in testis of the Nile tilapia *Oreochromis niloticus*. *Fish. Sci.*, **68(4)**, 830-837.
- Mohri M. and Nishida T., 2002: Considerationon horizontal and vertical distribution of adult yellowfin tuna (*Thunnus albacares*) in the Indian Ocean based on the Japanese tuna longline fisheries information. *La Mer*, **40**: 29-39.
- Mori K., Iida N., Nishzawa T., Arimoto M., and Nakajima K., 2002: Properties of viral hemorrhagic septicemia virus (VHSV) isolated from Japanese flounder *Paralichthys olivaceus*. *Fish Pathol.*, **37(4)**, 169-174.
- 森 賢, 中村好和, 2003: 黒潮親潮移行域周辺におけるスルメイカ幼体の分布. 日水誌, **69(1)**, 21-27.
- 森 賢, 木下貴裕, 佐々千由紀, 小西芳信, 2003: 黒潮周辺海域におけるスルメイカ冬季発生群の産卵海域と輸送経路. 月刊海洋, 号外31, 106-110.
- 森 信幸, 坪田幸雄, 福田光男, 丸山修治, 岸 哲也, 三宮芳明, 2002: 人工産卵基質による沿岸構造物への魚類産卵場機能の付加. 海岸工学論文集, **49**, 1336-1340.
- 森永健司, 吉村 拓, 2002: イセエビ属フィロソーマ幼生出現域の海洋環境. 月刊海洋, 号外31, 135-142.
- Morita K. and Morita A., 2002: Rule of age and size at maturity: individual variation in the maturation history of resident white-spotted charr. *J. Fish Biol.*, **61(5)**, 1230-1238.
- Morita T., 2002: Characteristics of  $\alpha$ -actin from the deep-sea fish *Coryphaenoides yaquinae*. *Fish. Sci.*, **68 (Suppl. )**, 1309-1310.
- 森実庸男, 山下浩史, 藤田慶之, 川上秀昌, 越智 僕, 前野幸男, 釜石 隆, 伊東尚史, 栗田 潤, 中島員洋, 芦田 勝朗, 2002: 血リンパ接種による軟 体部の赤変化を伴うアコヤガイ疾病の再現. 魚病研究, **37(3)**, 149-151.
- \*Munehara H., Nagai N., and Narimatsu Y., 2002: Isolation and characterization of microsatellite markers from the *Hoploptychus dybowskii* (Pisces, Gasterosteiformes). *Mol. Ecol. Notes*, **2**, 70-71
- Murano M. and Hanamura Y., 2002: A new species of *Heteromysis* (Crustacea, Mysida, Mysidae) from Japan. *Plankton Biol. Ecol.*, **49**, 75-80.
- 村田昌一, 2002: 我が国の海藻類の生産と利用. *Techno Innovation*, **12(1)**, 26-29.
- Murata Y., Sakaguchi M., Ninomiya Y., Sata N. U., and Oohara I., 2002: Taste quality and thresholds of a novel sulfur-containing amino acid, pulcherrimine in humans and mice. *Fish. Sci.*, **68(Suppl. )**, 1390-1393.
- 村田裕子, 金庭正樹, 大原一郎, 大村裕治, 岡崎恵美子, 2002: 飼育水温の変化が魚肉のエキス成分に及ぼす影響. 日本味と匂学会誌, **9(3)**, 745-746.
- Nagai S., Matsuyama Y., Takayama H., and Kotani Y., 2002: Morphology of *Polykrikos kofoidii* and *P. schwartzii* (Dinophyceae, Polykrikaceae) cysts obtained in culture. *Phycologia*, **41(4)**, 319-327.
- 永井達樹, 2002: 富栄養化からみた瀬戸内海の移り変わりと漁業資源の回復. 水産海洋研究, **66**, 190-191.
- Nagasaki K., Tarutani K., Tomaru Y., Itakura S., Tamai K., Yamanaka S., Tanabe H., Katanozaka N., Shirai Y., and Yamaguchi M., 2002: Possible use of viruses as a microbiological agent against harmful algal blooms. *Fish. Sci.*, **68(Suppl. )**, 497-500.
- Nakagawa Y., Endo Y., and Taki K., 2002: Contributions of heterotrophic and autotrophic prey to the diet of euphausiid, *Euphausia pacifica* in the coastal waters off northeastern Japan. *Polar Biosci.*, **15**, 52-65.
- Nakajima H., Takatsuki Y., Mizuno K., Takeuchi K. and Shikama N., 2002: Data communication status of the ARGO floats. *ARGO Technical Report FY2001, JAMSTEC*, pp.78-87.
- Nakajima K., Itoh N., Kurita J., Kawakami H., and Itano K., 2002: Effectiveness of a vaccine against red sea bream iridoviral disease in various cultured marine fish under laboratory conditions. *Fish Pathol.*, **37(2)**, 90-91.
- Nakamura M., Nagoya H., and Hirai T., 2002: Nonylphenol induces complete feminization of the gonad in genetically controlled all-male amago salmon. *Fish. Sci.*, **68(6)**, 1387-1389.
- 中村孝幸, 高木伸雄, 中山哲嚴, 河野 徹, 2002: ピストンモード波浪共振を利用する低反射・低透過構造の海水交換型防波堤の開発. 海岸工学論文集, **49**, 661-665.
- 中村孝幸, 高木伸雄, 中山哲嚴, 久保勝太, 飯干富広, 2002: 平面波浪場における垂下版式反射波低減工の効果について. 海岸工学論文集, **49**, 666-670.

- Nakamura T., Maruyama T., and Watanabe S., 2002: Residency and movement of stream-dwelling Japanese charr, *Salvelinus leucomaenoides*, in a central Japanese mountain stream. *Ecol. Freshwater Fish*, **11**, 150-157.
- 中村義治, 金網紀久恵, 磯野良介, 三村信男, 2002: 貝類の生物機能と水域環境への影響に関する全国評価. 海岸工学論文集, **49**, 1371-1375.
- 中村義治, 寺澤知彦, 中村幹男, 山下俊彦, 青木伸一, 2002: 物質循環スペクトルによる汽水湖生態系の健康度評価. 海岸工学論文集, **49**, 1446-1450.
- Nakanishi T., Kiryuu I., and Ototake M., 2002: Development of a new vaccine delivery method for fish: percutaneous administration by immersion with application of a multiple puncture instrument. *Vaccine*, **20**, 3764-3769.
- Nakata K., 2003: Interannual changes of the winter to early spring biomass and composition of mesozooplankton in the Kuroshio region in relation to climatic factors. *J. Oceanogra.*, **59(2)**, 225-234.
- Nakayama A., Yano Y., Yoshida K., Ishihara K., and Saito H., 2002: Characteristic and pressure of bacteria isolated from intestinal contents of deep-sea fish retrieved from the abyssal zone. *Fish. Sci.*, **68(Suppl.)**, 647-648.
- Nakayasu C., Yoshinaga T. and Kumagai A., 2002: Hematology of anemia experimentally induced by repeated bleedings in Japanese flounder anemia recently prevailing in Japan. *Fish. Pathol.*, **37**, 125-130.
- Nakayasu C., Tsutsumi N., Yoshitomi T., Yoshinaga T., and Kumagai A., 2003: Identification of Japanese flounder leucocyte population involved in the host response to *Neoheterobothrium hirame*. *Fish Pathol.*, **38(1)**, 9-14.
- Nanami A. and Nishihara M., 2002: The structures and dynamics of coral reef fish communities in an Okinawan coral reef. the effects of coral-based habitat structures at sites with rocky and sandy sea bottoms. *Environ. Biol. Fish.*, **63**, 353-372.
- \*Narimatsu Y., and Munehara H., 2001: Territoriality, egg desertion and mating success of a paternal care fish, *Hypoptichus dybowskii* (*Gasterosteiformes*). *Behaviour*, **135**, 85-96.
- 梨田一也, 本多仁, 阪地英男, 木村量, 2003: 足摺岬周辺及び土佐湾中央部海域で漁獲されたゴマサバの年齢形質としての耳石の有効性. 黒潮の資源海洋研究, **4**, 5-9.
- 新村陽子, 内村真之, 薄浩則, 吉川浩二, 吉田吾郎, 寺脇利信, 2002: 広島湾の主要海藻群落の外縁部における水環境と光透過率. 水産工学, **40**, 7-14.
- 新村陽子, 田口哲, 2002: 海氷と海水中の植物プランクトン. 月刊海洋, 号外**30**, 116-125.
- Nikaido M., Kawakami A., Tsuda A., Furuya-kiyoki M., Takeda H., and Araki, K., 2002: Tbx24, encoding a T-box protein, is mutated in the zebrafish somite-segmentation mutant. *Nature Genetics*, **31(2)**, 195-199.
- Nishida T., Meaden G. J., and Booth A. J., 2002: Spatial fish resources analyses using GIS (Geographical Information Systems): Current situation and prospects. *Fish. Sci.*, **68(Suppl.)**, 109-112.
- Nishi K., Chikae M., Hatano Y., Mizukami H., Sakakibara R., Yamashita M., and Tamiya E., 2002: Development and application of a monoclonal antibody-based sandwich ELISA for quantification of Japanese medaka (*Oryzias latipes*) vitellogenin. *Comp. Biochem. Physiol.*, **132**, 161-169.
- Nishimura A., Hamatsu T., Yabuki K., and Shida O., 2002: Recruitment fluctuations and biological responses of walleye pollock in the Pacific coast of Hokkaido. *Fish. Sci.*, **68(Suppl.)**, 206-209.
- 西村一彦, 鈴木敏之, 板橋豊, 2002: 蛍光検出HPLCによる乳及び乳製品中の遊離脂肪酸の定量. 食品衛生学雑誌, **43(4)**, 230-233.
- Niwa Y., Nakazawa A., Margulies D., Scholey V. P., Wexler J. B., and Chow S., 2003: Genetic monitoring for spawning ecology of captive yellowfin tuna (*Thunnus albacares*) using mitochondrial DNA variation. *Aquaculture*, **218**, 387-395.
- 野原健司, 岡村寛, 中立元樹, 平松一彦, 鈴木伸明, 岡崎誠, 張成年, 2003: メカジキ (*Xiphias gladius*) の吻内部に見られる2型の生物学的検討及び大西洋南北系群間の遺伝的差異. 水研センター研報, **7**, 1-13.
- Ogata H. Y., Oku H., and Murai T., 2002: Growth feed efficiency and feed intake of offspring from selected

- and wild Japanese flounder (*Paralichthys olivaceus*). *Aquaculture*, **211**, 183-193.
- Ogata H. Y., Oku H., and Murai T., 2002: Growth performance and macronutrient retention of offspring from wild and selected red sea bream (*Pagrus major*). *Aquaculture*, **206**, 279-287.
- Ogyri K., Mtsumoto Y., Yamada M., Saito Y., and Iseki K., 2002: Sediment accumulation rates and bugets of depositing particles of the East Chaina Sea. *Deep-sea Res.*, **50**, 513-528.
- Oh S. J., Yamamoto T., Kataoka Y., Matsuda O., Matsuyama Y., Kotani Y., 2002: Utilization of dissolved organic phosphorus by the two toxic dinoflagellates, *Alexandrium tamarens*e and *Gymnodinium catenatum* (Dinophyceae). *Fish. Sci.*, **68(2)**, 416-424.
- Ohizumi H., 2002: Dietary studies of toothed whales: A review of technical issues and new topics. *Fish. Sci.*, **68(Suppl. )**, 264-267.
- Ohizumi H., Isoda T., Kishiro T., and Kato H., 2003: Feeding habits of Baird's beaked whale, *Berardius bairdii*, in the western North Pacific and Sea of Okhotsk off Japan. *Fish. Sci.*, **69(1)**, 11-20.
- Ohizumi H. and Kishiro T., 2003: Stomach contents of a Cuvier's beaked whale (*Ziphius cavirostris*) stranded on the central Pacific coast of Japan. *Aquatic Mammals*, **29**, 99-103.
- Ohtsuka S., Hanamura Y., and Kase T., 2002: A new species of *Theti spelecaris* (Crustacea, Peracarida) from submarine cave on Grand Cayman Island. *Zool. Sci.*, **19**, 611-624.
- Oikawa H., Fujita T., Satomi M., Suzuki T., Kotani Y., and Yano Y., 2002: Accumulation of paralytic shellfish poisoning toxins in the edible shore crab *Telmessus acutidens*. *Toxicon*, **40(11)**, 1593-1599.
- Oikawa H., Stomi M., and Yano Y., 2003: Accumulation of paralytic shellfish poisonong toxins in the edibleshore crab *telmessus*. *Toxicon*, **40(11)**, 1593-1599.
- Oka E., Izawa K., Inoue A., Ando K., Shikama N., Mizuno K., Suehiro K., and Takeuchi K., 2002: Is retrieve of Argo floats possible?. *JAMSTECR*, **46**, 147-155.
- 岡本 誠 , 井田 斎 , 杉崎宏哉 , 栗田 豊 , 2002: 本州東方沖から得られたイレズミコンニャクアジの仔稚魚 . 魚類学雑誌 , **49(2)**, 97-102.
- Okamura H., Yatsu A., and Hiramatsu K., 2003: Fisheries management based on ecosystem models-A case study using Ecopath and Ecosim-. *Fish. Sci.*, **68(Suppl. )**, 154-157.
- 岡内正典 , 2002: 海産魚介類の初期餌料用微細藻類の大量培養技術の開発 . 日水誌 , **68(5)**, 625-628.
- 岡崎恵美子 , 山下由美子 , 大村裕治 , 2002: 高速攪拌による魚肉すり身中の魚油の微粒化と加熱ゲル形成能の向上 . 日水誌 , **68(4)**, 547-553.
- 岡崎恵美子 , 夜久俊治 , 福田 裕 , 酒井 徹 , 南部正一 , 2002: 凍結貯蔵中の乳化すり身の解乳化に及ぼすポリオール類の抑制効果 . 日水誌 , **68(5)**, 701-705.
- Okazaki T., Jeon S.-R., and Kitagawa T., 2002: Genetic differentiation of piscivorous chub (genus *Opsariichthys*) in Japan, Korea and Russia. *Zool. Sci.*, **19(5)**, 601-610.
- Oku H., Ogata H., and Liang X. F., 2002: Organization of the lipoprotein lipase gene of red sea bream *Pagrus major*. *Comp. Biochem. Physiol. B*, **131**, 775-785.
- Oku H., Liang X. F., and Ogata H. Y., 2002: Lipoprotein lipase gene in red sea bream. *Fish. Sci.*, **68(Suppl. )**, 1194-1197.
- Okumura T., Akamatsu T., Novarini N., and Yan H. Y., 2002: Analyses of small tank acoustics: Empirical and theoretical approaches. *Bioacoustics*, **12**, 330-332.
- 奥村都誓司 , 升也利一 , 高尾芳三 , 澤田浩一 , 2002: 水中生物の音響散乱への境界要素法の応用 . 海洋音響学会誌 , **30** , 58-65.
- Okumura T., Masuya T., Takao Y., and Sawada K., 2003: Scattering model by an arbitrary shape body. *Proc. ICES Symp.*, **60**, 563-570.
- Okuzawa K., 2003: Puberty in teleosts *Fish. Physiol. Biochem.*, **26**, 31-41.
- Ookubo K., Yamano K., Qin Q.W., Aoyagi K., and Ototake M., 2002: Ubiquitin genes in rainbow trout(*Oncorhynchus mykiss*). *Fish Shellfish Immunol.*, **12(4)**, 335-351.
- Ookubo K., Yoshiura Y., Amano K., Suetake H., and Aida K., 2003: The GnRH system in teleosts. *Aquatic*

- Genomics*, 244-262.
- Oomaki M., Saeki K., Kuroda K., Kenmochi S., 2002: Sheet pile mooring wharfs in soft ground: Centrifugal loading tests. Physical modelling in geotechnics: *ICPMG 2002*, 859-864.
- Oomaki M., Saeki K., and Takaki N., 2002: Improvement of marine environments by dredging and creation of marine products areas. *30th PIANC-AIPCN Congress*, pp.166-178.
- Oomaki M., Saeki K., and Takaki N., 2002: Development of an economical sheet pile mooring wharf. *30th PIANC-AIPCN Congress*, pp.461-467.
- Ohshima, S., Yoda, M., and Hiyama, Y., 2002: Attempt study of reproductive characteristics in two scombridae (*Scomber japonicus* and *Scomber australasicus*) in the East China Sea. *Proc. Process Dynamics Northwestern Pacific Ecosystems*, pp.15.
- Ohshima, S., 2002: Spatial distribution and biomass of pelagic fish in the East China Sea. *Proc. 6th ICES Symp. on Acoustics in Fisheries and Aquatic Ecology*, 87-88.
- 大関芳沖, 木村 量, 2002: 卵稚仔プランクトンのモニタリング (1)日本周辺海域の卵稚仔プランクトンモニタリング. *水産海洋研究*, **66**(4), 262-264.
- 大関芳沖, 2002: 日本周辺海域の卵稚仔プランクトンモニタリング. *月刊海洋*, **34**(11), 782-787.
- 大関芳沖, 2002: 黒潮親潮移行域を中心とするサンマ仔稚魚の分布・輸送と加入量変動. *月刊海洋*, 号外**31**, 178-183.
- Osako K., Yamaguchi A., Kurokawa T., Kuwahara K., Saito H., and Nozaki Y., 2002: Chemical components and body color of horse mackerel caught in different areas. *Fish. Sci.*, **68**(3), 587-594.
- 乙竹 充, Dijkstra J.M., 桐生郁也, 吉浦康寿, 藤原篤志, 2002: ニジマスMHC遺伝子の多様性及び機能の解析. *水産育種*, **32**, 9-16.
- Park, Y. C., Yoda, M., and Hiyama, Y., 2002: Stock assessment for swordtip squid, *Loligo edulis*, in the East China Sea and the southwest sea of Japan. *Fish. Sci.*, **68**(Suppl. ), 89-92.
- Qin Q.W., Otake M., Nagoya H., and Nakanishi T., 2002: Graft-versus-host reaction (GVHR) in clonal amago salmon, *Oncorhynchus rhodurus*. *Veter. Immunol. Immunopathol.* **89**, 83-89.
- Rhodes L., Suzuki T., Adamson J., and Mountfort D., 2001: Esterified okadaic acid in New Zealand strains of *Prorocentrum lima*. *Proc. 9th Intl. Conf. Harmful Algal Blooms*, 364-366.
- Saeki K., Takaki N., Akeda S., 2003: Evaluation and improvement of working environment at fishing ports. *PACON 2002*, 353-361.
- 佐伯公康, 穴井美緒, 朴 賢哲, 明田定満, 高木伸雄, 近藤健雄, 2002: 漁港における就労環境の評価・分析手法の構築. *海岸工学論文集*, **49**, 951-955.
- Saito H., Tsuda A., and Kasai H., 2002: Nutrient and plankton dynamics in the Oyashio region of the western subarctic Pacific. *Deep-Sea Res. II*, **49**: 5463-5486.
- Saito H., Kotani Y., Keriko J. M., Xue C., Taki K., Ishihara K., Ueda K., and Miyata S., 2002: High levels of n-3 polyunsaturated fatty acids in *Euphausia pacifica* and its role as a source of docosahexaenoic and icosapentaenoic acids for higher trophic levels. *Mar. Chem.*, **78**, 9-28.
- 齊藤宏明, 2002: 流体中の粒子遭遇理論を用いた動物プランクトン摂餌に関する研究. *日本プランクトン学会報*, **49**, 46-51.
- Saitoh H. and Yoshida K., 2002: Utilization of the lipids of the deep-sea grenadier: the high content of ceramide derivatives in the internal organ lipids of genus *Coryphaenoides*. *Fish. Sci.*, **68**(Suppl. ), 1426-1429.
- Saitoh H. and Taguchi S., 2003: Influence of UV-B radiation on hatching success of marine copepod *Paracalanus parvus* s. l. *J. Exp. Mar. Biol. Ecol.*, **282**, 135-147.
- Saitoh K., Miya M., Inoue J., Ishiguro N., and Nishida M., 2003: Mitochondrial genomics of ostariophysan fishes: perspectives on phylogeny and biogeography. *J. Molecular Evol.*, **56**, 464-472.
- 斎藤憲治, 2003: 農業土木技術者のための生き物調査(その7)－淡水魚調査法－. *農業土木学会誌*, **71**(3), 241-245.
- 齋藤暢宏, 梶原直人, 2002: 日本海新潟沿岸域から採集された大型アミ類2種(甲殻綱・アミ目・ロフオガスター科). *日本生物地理学会報*, **57**, 19-30.

- Sakai M. and Ishida K. 2002: The Pacific salmon ranching in the southern Chile: Evaluation of a project-type technical cooperation by JICA. *Fish. Sci.*, **68(Suppl. )**, 1952-1955.
- 阪地英男, 2003: 土佐湾におけるクルマエビ科エビ類の資源生物学的研究. 水研センター研報, **6**, 73-127.
- Sakano H. and Ueda H., 2002: Size-structured interaction in two zooplanktivorous fishes related to a rapid reduction of zooplankton biomass in a large oligotrophic lake. *Proceedings of the VIII INTECOL International Congress of Ecology*, 2002, 230-231.
- Sakurai Y., Kiyofuji H., Saito S., Mori K., and Kinoshita T., 2002: Stock fluctuations of the Japanese common squid, *Todarodes pacificus*, related to recent climate changes. *Fish. Sci.*, **68(Suppl. )**, 226-229.
- 櫻井泰憲, 山本 潤, 木所英昭, 森 賢, 2003: 気候のレジームシフトに連動したスルメイカの資源変動. 月刊海洋, **35(2)**, 100-106.
- Sano M., Minagawa M., Tamaki M., Hayasibara T., and Shimizu H., 2002: Effect of water temperature on the spawning interval of the Kuruma prawn *Panaeus japonicus* after unilateral eyestalk ablation. *Suisanzosyoku*, **50(4)**, 433-434.
- Sano M., Minagawa M., and Nakajima K., 2002: Multiplication of Red Sea Bream Iridovirus (RSIV) in experimentally infected grouper. *Fish Pathol.*, **37(4)**, 163-168.
- Saruwatari T., Oohara I., and Kobayashi T., 2002: Salangid fishes: their past, present and future. *Fish. Sci.*, **68(Suppl. )**, 71-74.
- 佐々千由紀, 小西芳信, 2002: 東シナ海におけるマアジ仔稚の分布と輸送. 月刊海洋, 号外**31**, 92-98.
- Satomi M., Oikawa H., and Yano Y., 2003: Shewanella marintesting sp. nov., *Shewanella schlegeliana* sp. nov., and *Shewanella sairae* sp. nov., novel eicosapentaenoic-acid-producing marine bacteria isolated from sea-animal intestine. *J. Syst. Evol. Microbiol.* **53**, 491-499.
- 佐藤真彦, 工藤雄一, 北村章二, 岩田宗彦, 竹内康人, 2002: 母川回帰行動の嗅覚機構解明へのニューロエソロジー(神経行動学)的アプローチ: 自由遊泳中のサケ科魚類の嗅覚中枢活動のテレメトリ. 横浜市大論叢自然科学系列, **53(3)**, 53-80.
- Sawada K., Takao Y., Miyahana Y., and Kinacigil H. T., 2002: Introduction of the precise TS measurement for fisheries acoustics. *Turkish J. Veteri. Animal Sci.*, **26**, 209-214.
- Sawada K. and Furusawa M., 2002: Computer Simulation study in situ target strength estimation method. Proceedings of International Symposium on Advanced Techniques of Sampling Gear and Acoustical Surveys for Estimation of Fish Abundance and Behavior (ACOUSTGEARE 2000), pp.11-16.
- Sekino M., Hara M., and Taniguchi N., 2002: Loss of microsatellite and mitochondrial DNA variation in hatchery strains of Japanese flounder *Paralichthys olivaceus*. *Aquaculture*, **213(1-4)**, 101-122.
- Sekino M., Hara M., and Takahashi, H., 2002: Molecular cloning of microsatellite DNA from marine organisms. *Tech. Rep. Hokkaido Natl. Fish. Res. Inst.*, **5**, 87-90.
- Sekino M., Saitoh K., Yamada T., Kumagai A., Hara M., and Yamashita Y., 2003: Microsatellite-based pedigree tracing in a Japanese flounder *Paralichthys olivaceus* hatchery strain: implications for hatchery management related to stock enhancement program. *Aquaculture*, **221**, 255-263.
- Sekino M., Hamaguchi M., Aranishi F., and Okoshi K., 2003: Development of novel microsatellite DNA markers from the Pacific oyster *Crassostrea gigas*. *Mar. Biotechnol.*, **5(3)**, 227-233.
- Sekino M., Hara M., and Taniguchi N., 2003: Genetic diversity within and between hatchery strains of Japanese flounder *Paralichthys olivaceus*. *UJNR Technical Report*, **30**, 43-50.
- 志田 修, 西村 明, 2002: 秋季の道東太平洋海域に分布するスケトウダラ0歳魚の孵化日組成と産卵群の関連. 水産海洋研究, **66(4)**, 232-238.
- Shibuno, T., Hashimoto, K., Abe, O., Takada, Y., and Kawasaki, H., 2002: Disturbed coral reefs and the effects upon the structure of fish communities at Ishigaki Island, Japan. *Fish. Sci.*, **68(Suppl. )**, 139-142.
- Shibuno, T., 2002: Study on the Disturbance and Restoration of Coral Reefs (Abstract of the Interim Report). *Global Environ. Res. Jpn.* 2001., 162-166.
- Shigehiro R., Aguilar G. D., Kuroda K., Kawai A., Matsuda A., 2002: Characteristics of maneuvering

- motions of Philippine Outrigger Craft. *J. Kansai Soc. Architects Japan*, **238**, 113-120.
- Shimizu A. and Yamashita M., 2002: Purification of mummichog (*Fundulus heteroclitus*) gonadotropins and their subunits, using an immunochemical assay with antisera raised against synthetic peptides. *Gen. Comp. Endocrinol.*, **125**, 79-91.
- Shimizu Y., Yasuda I., Okuda K., Hanada K., and Ito S., 2003: ADCP-referenced Kuroshio and Oyashio water transports for the North Pacific Intermediate Water formation. *J. Phys. Oceanogr.*, **33**(1), 220-233.
- 島 隆夫, 小西浩司, 古板博文, 山本剛史, 鈴木伸洋, 2002: 普及型自動給餌機を用いた自発摂餌型給餌装置の製作 . 水産増殖 , **50**(4), 441-442.
- 島谷 学, 中瀬浩太, 岩本裕之, 中山哲嚴, 月館真理雄, 星野高士, 内山雄介, 灘岡和夫, 2002: 興津海岸におけるアマモ分布条件について . 海岸工学論文集 , **49**, 1161-1165.
- Shiomoto A., Saitoh S., Imai K., Toratani M., Ishida Y., and Sasaoka K., 2002: Interannual variation in phytoplankton biomass in the Bering Sea basin in the 1990s. *Prog. Oceanogr.*, **55**(1-2), 147-164.
- Smith P. J., McMillan P. J., Bull B., McVeagh S. M., Gaffney P. M., and Chow S., 2002: Genetic and meristic variation in black and smooth oreos in the New Zealand Exclusive Economic Zone. *New Zealand J. Mar. Freshwater Res.*, **36**, 737-750.
- Srivastava A.S., Kurokawa T., and Suzuki T., 2002: mRNA expression of pancreatic enzyme precursors and estimation of protein digestibility in first feeding larvae of the Japanese flounder, *Paralichthys olivaceus*. *Comp. Biochem. Physiol.*, **A132**, 629-635.
- Suda M., Kishida T., and Akamine T., 2002: Recruitment of Japanese sardine, *Sardinops melanostictus*, off the Pacific coast of Japan during 1991-1995 estimated using a spatial numerical model. *Fish. Sci.*, **68**(Suppl. ), 439-440.
- Suda M. and Kishida T., 2003: A spatial model of population dynamics of the early life stages of Japanese sardine, *Sardinops melanostictus*, off the pacific coast of Japan. *Fish. Oceanogr.*, **12**(2), 85-99.
- 巣山 哲, 2002: 北太平洋におけるサンマ Cololabis saira (Brevoort) の年齢, 成長および成熟に関する研究 . 水研センター研報 , **5**, 68-113.
- \*Suzuki T., Oota H., and Yamazaki M., 2001: Dinophysistoxin-1 and esterified dinophysistoxin-1 in the mussel *Mytilus galloprovincialis* fed on the toxic dinoflagellate. *Proc. 9th Int'l. Conf. Harmful Algal Blooms*, 367-370.
- Suzuki T., Yamazaki M., and Kamiyama T., 2002: The Japanese shellfish toxin monitoring system. *Proc. 9th Int'l. Symp. Toxic Micro-organisms*, pp.148-157.
- Suzuki T., Beuzenberg V., Mackenzie L., and Quiliam M. A., 2003: Liquid chromatography-mass spectrometry (LC-MS) of spiroketal stereoisomers of pectenotoxins and the analysis of novel pectenotoxin isomers in the toxic dinoflagellate *Dinophysis acuta* from New Zealand. *J. Chromatogr. A*, **992**, 141-150.
- Suzuki T., Srivastava A. S., Kurokawa T., 2002: A homologue of human placental protein, PP11, and mouse T cell-specific protein, Tcl-30, in the exocrine pancreas of a teleost (*Paralichthys olivaceus*). *Comp. Biochem. Physiol.*, **B133**, 325-329.
- Suzuki N., Suzuki T., Kurokawa T., 2002: Possible involvement of calcitonin gene-related peptide in seawater adaptation of flounder: Expression analysis of its receptor mRNA in the gill. *Fish. Sci.*, **68**(2), 425-429.
- Tada M., 2002: Econometric analysis of decreasing trend of bigeye tuna resources in the Indian Ocean. *Proc. Internl. Symp. Ocean Fish./China*, 60-72.
- 多田 稔, 有路昌彦, 2003: コンブとワカメの需給動向に関する経済分析 . 北日本漁業 , **31**, 56-66.
- Takada, Y., Kosuge, T., Nishihama, S., and Katoh, M., 2002: Vertical distributions of *Atactodea striata* and *Latona faba* (*Bivalvia*) on a subtropical sandy beach in Ishigaki Island, Japan. *Venus*, **61**, 203-213.
- \*Takada, Y. and Abe, O., 2002: Local variations in shootdensity, shoot morphology, and bite mark frequency of subtropical seagrasses in Japan. *Bull. Fish. Res. Agen.*, **4**, 1-10.
- Takada, Y., 2002: Activity patterns of herbivorous molluses on boulder shores. *Aquabiol.*, **143**, 484-490.
- Takahashi I., Nagao N., and Endo Y., 2002: Respiration of adult female *Calanus hyperboreus* (Copepoda)

- during spring in the North Water Polynya. *Polar Biosci.*, **15**, 45-51.
- Takahashi S. and Okazaki T., 2002: A new lentic form of the "yoshinobori" species complex, *Rhinogobius* spp. from Lake Biwa, Japan, compared with lake-river migrating *Rhinogobius* sp. OR. *Ichthyol. Res.*, **49**(4), 333-339.
- Takami H., Kawamura T., and Yamashita Y., 2002: Effects of delayed metamorphosis on larval competence, and post-larval survival and growth of abalone *Haliotis discus hannai*. *Aquaculture*, **213**(1-4), 311-322.
- 高田宜武, 2002: 転石海岸における藻食性貝類の周期的な摂餌活動, 海洋と生物, **143**, 484-490.
- Takatsuki Y., Ichikawa Y., Kobayashi T., Mizuno K. and Takeuchi K., 2002: Construction of the automated data processing and delayed-mode quality control system for profiling floats. *ARGO Technical Report FY2001*, JAMSTEC, pp. 88-99.
- 瀧 憲司, 壺 純子, 中川至純, 遠藤宜成, 2002: 道東及び常磐沿岸域におけるツノナシオキアミ*Euphausia pacifica* の胃内容物の特徴. 水産海洋研究, **66**(3), 155-163.
- Tamaki H., Tokuoka M., Nishijima W., Terawaki T., and Okada M., 2002: Deterioration of eel-grass, *Zostera marina* L., meadows by water pollution in Seto Inland Sea, Japan. *Mar. Pollut. Bull.*, **44**, 1253-1258.
- 玉置 仁, 西嶋 渉, 富永春江, 寺脇利信, 岩瀬晃盛, 岡田光正, 2002: 広島県沿岸域における透明度変化とアマモ面積増減との関係. 水環境学会誌, **25**, 151-156.
- 玉置泰司, 2002: 浮魚礁の経済効果 - 鹿児島県奄美大島海区を事例として -. 地域漁業研究, **42**(2), 103-118.
- 田中勝久, 豊川雅哉, 澤田知希, 柳澤豊重, 黒田伸郎, 2003: 土壤流出によるリン負荷の沿岸環境への影響. 沿岸海洋研究, **40**(2), 131-139.
- 田坂行男, 2003: 産地統合化時代の物流問題. 漁業経済研究, **47**(3), 29-43.
- Terasaki M., Itabashi Y., Suzuki T., and Nishimura K., 2002: An improved method for determining the composition of FFA in red tide flagellates by RP-HPLC with fluorescence detection. *J. Am. Oil Chem. Soc.*, **79**(12), 1181-1186.
- Terawaki T., Yoshikawa K., Yoshida G., Uchimura M., and Iseki K., 2002: Ecology and restoration techniques for Sargassum beds in Seto Inland Sea, Japan. *Mar. Pollut. Bull.*, **47**, 198-201.
- 寺脇利信, 吉川浩二, 吉田吾郎, 内村真之, 新井章吾, 2002: 南西日本の磯焼け海域における海底景観の特徴. 水産工学, **39**, 29-35.
- 寺脇利信, 吉見圭一郎, 2002: お掃除フリーの海藻栽培水槽の試み 4. ヤツマタモクとモズクの生長 海苔と海藻, **64**, 22-26.
- 寺脇利信, 吉川浩二, 吉田吾郎, 内村真之, 新村陽子, 2002: 広島湾における藻場の生態と回復技術. 水産海洋研究, **66**, 195-197.
- 寺脇利信, 新井章吾, 2002: 藻場の景観模式図9. 宮崎県門川湾乙島地先. 藻類, **50**, 21-23.
- 寺脇利信・新井章吾, 2002: 藻場景観模式図10. 新潟県佐渡島・真野湾二見地先. 藻類, **50**, 23-25.
- 寺脇利信, 新井章吾, 2002: 藻場の景観模式図11. 北海道厚岸郡浜中町藻散布地先の投石事業地. 藻類, **50**, 117-119.
- Tian Y., Akamine T., and Suda M., 2003: Variations in the abundance of Pacific saury (*Cololabis saira*) from the northwestern Pacific in relation to oceanic-climate changes. *Fish. Res.*, **60**(2), 439-454.
- Tohata K., Tokuda Y., Sakaguchi M., and Toyohara H., 2002: Immunological detection of translation products of type V/XI collagen a1 gene and their degradation products in red sea bream. *Fish. Sci.*, **68**(5), 1118-1123.
- 陶山典子, 輿石裕一, 須田有輔, 村井武四, 2003: 底質から見た有明海北部の海域区分とマクロベントスの分布. 水大校研報, **51**(4), 105-114.
- \*辻野 瞳, 吉田吾郎, 内田卓二, 2002: 宮島干潟におけるメイオベントス動物群-特に線虫類の現存量について-. 水研センター研報, **4**, 33-39.
- Tsujino M., Kamiyama T., Uchida T., Yamaguchi M., and Itakura S., 2002: Abundance and germination capability of resting cysts of *Alexandrium* spp. (Dinophyceae) from faecal pellets of macrobenthic organisms. *J. Exp. Mar. Biol. Ecol.*, **271**, 1-7.

- Tsukamoto Y., 2002: Leptocephalus larvae of *Pterothrius gissu* collected from the Kuroshio-Oyashio region of the western North Pacific, with comments on its metamorphosis. *Ichthyol. Res.*, **49**, 267-269.
- Tsukamoto Y., Yamada H., and Zenitani H., 2002: Microincrements of otoliths of the Japanese sand lance *Ammodytes personatus* during early life stages. *Fish. Sci.*, **68(5)**, 1158-1160.
- Tsunogai S., Iseki K., Kusakabe M., and Saito Y., 2002: Biochemiacal cycles in the East Chaina Sea: MASFLEX program. *Deep-sea Res.*, **50**, 321-326.
- Uchida D., Yamashita M., Kitano T., and Iguchi T., 2002: Oocyte apoptosis during the transition from ovary-like tissue to testes during sex differentiation of juvenile zebrafish. *J. Exp. Biol.*, **205**, 711-718.
- Uchida M. and Murata M., 2002: Fermentative preparation of single cell detritus from seaweed, *Undaria pinnatifida*, suitable as a replacement hatchery diet for unicellular algae. *Aquaculture*, **207**, 345-357.
- Uchida M., Maeda T., and Shiba T., 2002; Phylogenetic analysis of three marine bacteria that have an ability to decompose *Laminaria japonica*. *Fish. Sci.*, **68(3)**, 703-705.
- Uchida M., Numaguchi K., and Murata M., 2002: Development of "marine silage" with a scope for constructing aquaculture for bioremediation, not bioremediation for aquaculture. *Fish. Sci.*, **68(Suppl. )**, 859-862.
- 内田基晴, 2002: 海藻の発酵について. 日本乳酸菌学会誌, **13(2)**, 92-113.
- Uchikawa K., Yamamura O., Kitagawa D., and Sakurai Y., 2002: Diet of the mesopelagic fish *Notoscopelis japonicus* (Family:Myctophidae) associated with the continental slope off the Pacific coast of Honshu, Japan. *Fish. Sci.*, **68(5)**, 1034-1040.
- Uehara S. and Mitani T., 2002: Horizontal and diel vertical distribution of eggs and larvae of two clupeoid fish (*Etrumeus teres* and *Sardinops melanostictus*). *Fish. Sci.*, **68(Suppl. )**, 435-436.
- Ueno D., Iwata H., Tanabe S., Ikeda K., Koyama J., and Yamada H., 2002: Specific accumulation of persistent organochlorines in bluefin tuna collected from Japanese coastal waters. *Mar. Poll. Bull.*, **45**, 254-261.
- 上野成三, 永田良助, 山崎英治, 中山哲嚴, 2002: 赤潮・貧酸素をリアルタイムに観測する野見湾漁場環境情報システム (Nomi Bay watch system) の開発. 海岸工学論文集, **49**, 1531-1535.
- Umeda N., Iskandar B. H., Hashimoto H., Urano S., Matsuda A., 2002: Comparison of European and Asia trawlers-stability in seaways-. *Proc. 1st Asia Pacific Workshop Mar. Hydrodynamics*, pp.79-84.
- Umeda N., Urano S., A. Matsuda A., and Hashimoto H., 2002: Anti-broaching steering system-comparison between experiment and calculation-. *Proc. 1st Asia Pacific Workshop Mar. Hydrodynamics*, pp.85-90.
- Unuma T., Yamamoto T., Akiyama T., Shiraishi M., Ohta H., 2003: Quantitative changes in yolk protein and other components in the ovary and testis of the sea urchin *Pseudocentrotus depressus*. *J. Exp. Biol.* **206**, 365-372.
- Urushidani H., Shimizu A., Mori Y., and Iguchi T., 2002: Early estrogen exposure induces abnormal development of *Fundulus heteroclitus*. *J. Exp. Zool.*, **293(7)**, 693-702.
- Usuki H., Hamaguchi M., and Ishioka H., 2002: Effects of developmental stage, seawater concentration and rearing temperature on cryopreservation of Pacific oyster *Crassostrea gigas* larvae. *Fish. Sci.*, **68(4)**, 757-762.
- Wada K., 2002: Perspective of genetic improvement in the culture of pearl oyster. *Fish. Sci.*, **68(Suppl. )**, 726-733.
- 若林 輝, 中村智幸, 久保田仁志, 丸山 隆, 2002: 中禅寺湖流入河川におけるサケ科魚類3種の産卵生態. 魚類学雑誌, **49(2)**, 133-141.
- Watanabe C. and Nishida H., 2002: Development of assessment techniques for pelagic fish stocks: applications of daily egg production method and pelagic trawl in the Northwestern Pacific Ocean. *Fish. Sci.*, **68(Suppl. )**, 97-100.
- Watanabe H., Kawaguchi K. and Hayashi A., 2002: Feeding habits of juvenile surface migratory myctophid fishes (Family Myctophidae) in the Kuroshio region of the western North Pacific. *Mar. Ecol. Prog. Ser.*, **236**, 263-272.

- Watanabe H. and Kawaguchi K., 2003: Decadal change in abundance of surface migratory myctophid fishes in the Kuroshio region from 1957 to 1994. *Fish. Oceanogr.*, **12**, 1-12.
- 渡邊一功, 本多 仁, 渡邊朝生, 内山雅史, 八角直道, 松本育夫, 永嶋 宏, 小野寺光文, 小坂 淳, 2003: マイワシ豊漁期・不漁期における房総以北海域の来遊資源動向と海洋 条件の変化 . 黒潮の資源海洋研究 , **4**, 23-34.
- Watanabe T., 2002: Method to estimate the population density of the snow crab *Chionoecetes opilio* using a deep-sea video monitoring system on a towed slegge. *Jpn. Agri. Res. Quart.*, **36**, 51-57.
- 渡部俊広, 渡部一俊, 北川大二, 2002: ズワイガニ類とキチジに対するトロ - ル網の漁獲効率 . 東北底魚研究 , **22**, 32-33.
- Watanabe T., 2002: Development of the deep-sea video monitoring system on a towed slegge to estimate the population density of the snow crab *Chionoecetes opilio*. *Fish. Sci.*, **68(Suppl. )**, 101-104.
- Waters J. M., Saruwatari T., Kobayashi T., Oohara I., McDowall R. M., and Wallis G., 2002: Phylogenetic placement of Retropinnid fishes: Data set incongruence can be reduced by using asymmetric character state transformation costs. *Syst. Biol.*, **51(3)**, 432-449.
- Xia C., Kiryu I., Dijkstra, J.M., Azuma T., and Nakanishi. T., 2002: Differences in MHC class I genes between strains of rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.*, **12(4)**, 287-301.
- Yada T., Uchida K., Kajimura S., Azuma T., and Hirano T., 2002: Immunomodulatory effects of prolactin and growth hormone in the tilapia. *J. Endocrinol.*, **173**, 483-492.
- 八木 宏, 足立久美子, 2002: 那珂川河口海域における物質分布構造の季節変化 . 海岸工学論文集 , **49**, 1026-1030.
- 八木 宏, 足立久美子, 2002: 黒潮流路変動に伴う沿岸域への栄養塩流入現象 . 海岸工学論文集 , **49**, 1216-1220.
- 八木 宏, 中山哲嚴, 足立久美子, 2002: 黒潮流路変動と前線渦が鹿島灘の広域沿岸水に与える影響 . 土木学会論文集 , **719**, 81-91.
- Yamada H., Amano K., Okuzawa K., Tiba H., and Iwata M., 2002: Maturational changes in brain contents of salmon GnRH in rainbow trout as measured by a newly developed time-resolved fluoroimmunoassay. *Gen. Comp. Endocrinol.*, **126(2)**, 136-143.
- Yada T. and Nakanishi T., 2002: Interaction between endocrine and immune systems in fish. *Internl. Rev. Cytol.*, **220**, 35-92.
- Yada T., Moriyama S., Suzuki Y., Azuma T., Hirose S. and Naito N., 2002: Relationships between obesity and metabolic hormones in the cobalt variant of rainbow trout. *Gen. Comp. Endocrinol.*, **128**, 36-43.
- Yada T., Uchida K., Kajimura S., Azuma T., and Hirano T., 2002: Immunomodulatory effects of prolactin and growth hormone in the tilapia, *Oreochromis mossambicus*. *J. Endocrinol.*, **127(3)**, 223-231.
- 山田 久, 角埜 彰, 2003: 非有機スズ系代替船底防汚塗料の開発状況と水生生物に対する有害性評価 . 水研センター研報 , **6**, 56-73.
- Yamada Y., Yokoyama H., Ishihi Y., and Azeta M., 2003: Historical feeding analysis in fish farming based on carbon and nitrogen stable isotope ratio in sediment. *Fish. Sci.*, **69(1)**, 213-215.
- 山田佳裕, 丸山 敦, 石樋由香, 2002: 沿岸帯における炭素, 窒素安定同位体比研究の話題 . 陸水学雑誌 , **63(3)**, 261-267.
- Yamaguchi M., Itakura S., Nagasaki K., and Kotani Y., 2002: Distribution and abundance of resting cysts of the toxic *Alexandrium* spp. (Dinophyceae) in sediments of the western Seto Inland Sea, Japan. *Fish. Sci.*, **68(5)**, 1012-1019.
- Yamaguchi, S., Gen K., Okuzawa K., Matuyama M., Kagawa H., 2002: Effects of gonadal steroids on gonadotropin subunit genes expression in male red seabream (*Pagrus major*). *Fish. Sci.*, **68(Suppl. )**, 953-954.
- 山本 潤, 中山哲嚴, 時吉 学, 宮地健司2002: 野見湾における夏期の流況と水質変動に関する現地観測 . 海岸工学論文集 , **49**, 1086-1090.
- Yamamoto T., Oh S. J., and Kataoka Y., 2002: Effect of temperature, salinity and irradiance on the growth of the toxic dinoflagellate *Gymnodinium catenatum* (Dinophyceae) isolated from Hiroshima Bay, Japan. *Fish. Sci.*, **68(2)**, 356-363.

- Yamamoto T., Seike T., Hashimoto T., and Tarutani K., 2002: Modelling the population dynamics of the toxic dinoflagellate *Alexandrium tamarensense* in Hiroshima Bay, Japan. *J. Plankton Res.*, **24**, 33-47.
- 山本民次, 橋本俊也, 辻 けい子, 松田 治, 樽谷賢治, 2002: 1991~2000年の広島湾海水中における親生物元素の時空間的変動, 特に植物プランクトン態 C:N:P比のレッドフィールド比からの乖離. *沿岸海洋研究*, **39**, 163-169.
- Yamamura O., Honda S., Shida O., and Hamatsu T., 2002: Diets of walleye pollock *Theragra chalcogramma* in the Doto area, northern Japan: ontogenetic and seasonal variations. *Mar. Ecol. Prog. Ser.*, **238**, 187-198.
- Yamashita Y., Tominaga O., Takami H., and Yamada H., 2003: Comparison of growth, feeding and cortisol level in stone flounder *Platichthys bicoloratus* juveniles between estuarine and nearshore nursery grounds. *J. Fish Biol.*, **63**(3), 617-630.
- 山崎慎太郎, 日向野純也, 渡部俊広, 2002: 貝桁網によるチョウセンハマグリの足部損傷について. *日水誌*, **68**(3), 368-373.
- Yamamoto T., Hashimoto T., Tarutani K., and Kotani Y., 2002: Effects of winds, tides and river water runoff on the formation and disappearance of the *Alexandrium tamarensense* bloom in Hiroshima Bay, Japan. *Harmful Algae*, **3**, 301-312.
- Yamamoto T., Shima T., Furuita H., and Suzuki N., 2002: Influence of dietary fat level and whole body adiposity on voluntary energy intake by juvenile rainbow trout *Oncorhynchus mykiss* (Walbaum) under self-feeding conditions. *Aquaculture Res.*, **33**(9), 715-723.
- Yamamoto T., Shima T., Furuita H., and Suzuki N., 2002: Influence of feeding diets with and without fish meal by hand and by self-feeders on feed intake, growth and nutrient utilization of juvenile rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, **214**, 289-305.
- Yamashita M., 2002: Heat shock response in zebrafish embryo. *Fish. Sci.*, **68**(Suppl. ), 1029-1032.
- Yanagimoto T., Nishimura A., Mito K., Takao Y., and Williamson N., 2002: Interannual changes of biological properties of walleye pollock *Theragra chalcogramma* in the central Bering Sea. *Prog. Oceanogr.*, **55**, 195-208.
- 柳本 順, 北村 徹, 2002: mtDNAシトクロームb遺伝子による北海道周辺のタラ科3種の種判別. *日水誌*, **68**(6), 893-899.
- Yano, K. and Matsuura, K., 2002: A review of the genus *Oxynotus* (Squaliformes, Oxynotidae). *Bull. Natn. Sci. Mus., Tokyo. Ser.A*, **28**, 109-117.
- Yano, K., Matsuura, K., and Tsukada, O., 2002: Redescription of the rare squaloid shark *Oxynotus japonicus* from Suruga Bay and the Enshu-nada Sea, Japan. *Species Diversity*, **7**, 363-369.
- Yano, K. and Shimada, K. 2002: Diel movement behavior of the tiger shark, *Galeocerdo cuvier*, at the Miyako Islands, Japan using ultrasonic telemetry. *82nd Annual Meet. Am. Soc. Ichthyol. Herpetol.*, pp.311.
- Yano, K., Yamada, H., and Kosuge, T., 2002: Migration behavior of the Pacific bluefin tuna in the spawning areas of the Ryukyu Islands using pop-up satellite tagging and ultrasonic telemetry. *Abstracts for 82nd Annual Meeting of the American Society of Ichthyologist and Herpetologists*, pp.311-312.
- 矢野和成, 2002: 日本周辺に生息するサメ類の種類、生態、人間との関係. *海洋と生物*, **142**, 373-374.
- 矢野和成, 2002: サメの攻撃による人的被害と被害による社会現象. *海洋と生物*, **142**, 415-423.
- Yatsu A., Mitani T., Watanabe C., Nisida H., Kawabata淳., and Matsuda H., 2002: Current stock status and management of chub mackerel, *Scomber japonicus*, along the Pacific coast of Japan -an example of allowable biological catch determination-. *Fish. Sci.*, **68**(Suppl. ), 93-96.
- \*Yoshimura, T., Morinaga, K., Shirai, S., and Yamakawa, H., 2002: Palinurid phyllosoma larvae and their distribution in winter off the Pacific coast of Japan. *Fish. Sci.*, **68**(1), 194-197.
- Yoda, M. and Aoki, M., 2002: Comparative study of benthic and pelagic populations of *Bodotria similis* (Cumacea) from Izu Peninsula, southern Japan. *J. Crustacean Biol.*, **22**(3), 543-552.
- Yoda, M. and Yoneda, M., 2002: Assessment of spawning frequency and batch fecundity in the yellow sea

- bream, *Dentex tumifrons*. *Fish. Sci.*, **68(Suppl. )**, 443-444.
- Yoda, M., Ohshima, S., and Hiyama, M. 2002: Stock assessment of jack mackerel in the East China Sea and adjacent waters. *China Japan Korea Joint GLOBEC Symp.*, pp.14.
- 淀 太我 , 井口恵一朗 , 2003: 外来種コクチバスの河川内繁殖の確認 . 水産増殖 , **51(1)**, 31-34.
- Yokoyama H., 2002: Impact of fish and pearl farming on the benthic environments in Gokasho Bay: Evaluation from seasonal fluctuations of the macrobenthos. *Fish. Sci.*, **68(2)**, 258-268.
- Yokoyama H., Higano J., Adachi K., Ishihi Y., Yamada Y., and Pichitkul P., 2002: Evaluation of shrimp polyculture system in Thailand based on stable carbon and nitrogen isotope ratios. *Fish. Sci.*, **68(4)**, 745-750.
- 横山 寿 , 西村昭史 , 井上美佐 , 2002: 熊野灘沿岸の魚類養殖場におけるマクロベントス群集と堆積物に及ぼす養殖活動と地形の影響 . 水産海洋研究 , **66**, 133-141.
- 横山 寿 , 西村昭史 , 井上美佐 , 2002: マクロベントスの群集型を用いた魚類養殖場環境の評価 . 水産海洋研究 , **66**, 142-147.
- Yoshida H., 2002: Population structure of finless porpoise (*Neophocaena phocaenoides*) in coastal waters of Japan. *The Raffles Bull. Zool.*, **Suppl. 10**, 35-42.
- 吉田英可, 2002: 鯨類の資源量推定のための飛行機目視調査 - スナメリに対する調査 - . 海洋と生物, **24**, 335-340.
- Yoshimura, T., Morinaga, K., Shirai, S., and Kiyomoto, S., 2002: Environmental factors affecting larval and post-larval recruitment of the Japanese spiny lobster, *Panulirus japonicus*. *Fish. Sci.*, **68(suppl. )**, 194-197.
- 芳村康男 , 馬 寧 , 鈴木四郎 , 梶原善之 , 2002: バルジによる改造漁船船型の操縦性能 . 日本造船学会論文集 , **192**, 37-46.
- 塙本達也 , 2002: 藻食性魚類の漁獲・利用の事例 . 水産工学会 , **30(1)**: 37-40.
- 塙本達也 , 児玉正昭 , 2002: マフグに腹空内接種した炭素粒子の組織内分布 (短報) . 水産増殖 , **50(3)**, 381-382.