

Abstracts of Poster Presentations

List of Presentations

- 1: Marine microalga *Rhodomonas* sp. as a novel live food for bivalves (S. Yamamoto, M. Okauchi and T. Yoshimatsu)
- 2: Observation and monitoring of asari clam predators in a population-collapsed habitat using Field Server (N. Tezuka, N. Kajihara, T. Fukatsu, K. Oguri and T. Yamakita)
- 3: Seasonal variations of carbon and nitrogen stable isotope ratios in Manila clam, *Ruditapes philippinarum*, in Wajiro intertidal sand flat, Japan (M. Kodama, M. Yamada, S. Yamazaki and N. Ikeda)
- 4: Predator-prey relationship among 3 species (Manila clam, kuruma prawn, tiger puffer) for stock enhancement program at tidal flat in Japan (A. Yoshida, H. Yamane, H. Yamazaki, A. Itou, K. Sakiyama and Y. Sakakura)
- 5: Correlation between water nutrient level and catch of asari, *Ruditapes philippinarum*, in fishing grounds in Japan (M. Uchida, M. Tsujino, N. Tezuka, T. Shigeta, Y. Takada, K. Niwa and Y. Ishihi)
- 6: Long-term fluctuations in stock abundance of the asari clam (*Ruditapes philippinarum*) in a no-fishing area in the southern part of Ise Bay, Japan (K. Hanyu)
- 7: Effects of small clam abundance, rainfall and sediment on recruitment success of the asari clam (*Ruditapes philippinarum*) in a no-fishing area in the southern part of Ise Bay, Japan (K. Hanyu)
- 8: Sampling design for abundance estimation of the asari clam (*Ruditapes philippinarum*) in Ise Bay, Japan (K. Hanyu, H. Kokubu, N. Hata, T. Mizuno, T. Inoue, T. Kudo, N. Hasegawa, Y. Ishihi, S. Watanabe, Y. Fujioka, J. Higano, R. Nambu and H. Kuwahara)
- 9: Discovery of a spawning ground of the asari clam (*Ruditapes philippinarum*) in Ise Bay, Japan (K. Hanyu, H. Kokubu, N. Hata, T. Mizuno, T. Inoue, Y. Tanaka, T. Kudo, N. Hasegawa, Y. Ishihi, S. Watanabe, Y. Fujioka, J. Higano, R. Nambu and H. Kuwahara)
- 10: Ontogenetic changes in ingestion and digestion abilities against diatom cells of the Manila clam (S. Houki, T. Kawamura, N. Ogawa, N.I. Won and Y. Watanabe)
- 11: Reproductive biology of Manila clam *Ruditapes philippinarum* in the coastal waters along eastern Hokkaido, Japan (Y. Konda and M. Sasaki)
- 12: Food environment for bivalves around *Zostera marina* beds (H. Kokubu, H. Hanyu and N. Hata)
- 13: Manila clam, *Ruditapes philippinarum*, culture using *Pyropia* spheroplast in closed recirculation system (N. Arisman, M. N. D. Khan, A. M. Shahabuddin, T. Yoshimatsu, T. Araki)
- 14: Comparison of types of cages and substrates in suspended culture of the Manila clam, *Ruditapes philippinarum* (N. Hata, N. Hasegawa, T. Mizuno, Y. Fujioka, Y. Ishihi, S. Watanabe, D. Asao, M. Yamaguchi, H. Imai, K. Morita and J. Higano)
- 15: Effects of feeding regimes on oocyte quality of Manila clam *Ruditapes philippinarum* (M. Awaji, T. Matsumoto and J. Higano)
- 16: Low-salinity endurances of the penshell *Atrina* sp. and clam *Trapezium liratum* (T. Kurihara, S. Nakano, Y. Matsuyama, K. Hashimoto, A. Ito and M. Kanematsu)
- 17: Effect of netting on survival of Manila clam *Ruditapes philippinarum* in an artificial tideland in Okayama, Japan (K. Izumikawa, T. Mototani, F. Murayama, J. Sato, A. Nishimoto, H. Yazamazaki, K. Sakiyama, and A. Ito)
- 18: Effects of water temperature on the expression of heat shock protein (HSP) 22, 40, and 70 genes in Manila clam (*Ruditapes philippinarum*) larvae (D. Ojima, M. Kanematsu, A. Nishimoto, A. Ito, H. Yamazaki, K. Asami and K. Sakiyama)
- 19: Spatiotemporal population structure of the asari clam *Ruditapes philippinarum* juvenile in Ise Bay inferred by microsatellite marker analysis

- (N. Sano, N. Hasegawa, K. Hanyu, D. Miyawaki, A. Komaru and H. Kuwahara)
- 20: Present situation of the asari clam *Ruditapes philippinarum* resources in Mikawa Bay, Japan (D. Miyawaki, Y. Murauchi, M. Hibino and S. Okamoto)
 - 21: Seasonal dynamics of the parasitic larvae and free-living adults of the sea spider *Nymphonella tapetis* infecting the asari *Ruditapes philippinarum* in the eastern coast of Chita Peninsula (Y. Murauchi, S. Okamoto, A. Hirai, D. Miyawaki, N. Yamamoto and M. Hibino)
 - 22: Shellfish production and new approaches in Turkey (S. Sedar)
 - 23: Variations of glycogen, protein and lipid contents depend on size of *Ruditapes decussatus* in Bostanli Seashore Izmir Bay, Aegean Sea, Turkey (S. Serdar, H. Pulat and G. Gungor)
 - 24: Use of *Pyropia* spheroplasts as a live food substitute for culturing Manila clam, *Ruditapes philippinarum* (A.M. Shahabuddin, M.N.D. Khan, N. Arisman, D. Saha, T. Yoshimatsu and T. Araki)
 - 25: Main ecological processes for the determination of *Ruditapes philippinarum* population size on an extensive intertidal sandflat in Ariake Sound, Kyushu, Japan (S. Takeuchi and A. Tamaki)
 - 26: Age determination of Asari *Ruditapes philippinarum* using growth rings on shell surface and section in Mutsu Bay, northern Japan (D. Sugiura and N. Kikuya)
 - 27: Abundance of planktonic larvae of the asari clam *Ruditapes philippinarum* around fishing grounds in Mikawa Bay (N. Kuroda, S. Okamoto, T. Matsumura and M. Hamaguchi)
 - 28: Population status of Manila clam, *Ruditapes philippinarum*, and physical environmental conditions in the Natori River estuary in northeastern Japan after the Great Tsunami (K. Ito, H. Yoshida, Y. Morioka, A. Katayama, D. Takahashi, M. Shoji, K. Sasaki and S. Katayama)
 - 29: Dynamics of the sea spider parasitism on asari in Matsukawaura Lagoon, Fukushima, Japan (T. Tomiyama, T. Iwasaki, K. Wakui, K. Yamada, M. Tamaoki and K. Miyazaki)
 - 30: Life history and distribution of a sea spider, *Nymphonella tapetis*, parasitic on the Manila clam, *Ruditapes philippinarum*, in Tokyo Bay, Japan (K. Yamada, K. Miyazaki, S. Chow, T. Yamamoto, T. Tomiyama, T. Yoshinaga, Y. Miyama, M. Tamaoki and M. Toba)
 - 31: Distribution, morphology and mode of life in the sea spider *Nymphonella* spp. (Arthropoda, Pycnogonida): Potential harmful endoparasitism on asari outside Japan (K. Miyazaki, T. Tomiyama, K. Yamada and M. Tamaoki)
 - 32: Larval and juvenile dynamics of the Manila clam *Ruditapes philippinarum* in northeastern Japan (H. Abe, M. Hamaguchi, N. Kajihara, Y. Taniai, A. Oshino, T. Iwasaki, T. Wada and T. Kamiyama)
 - 33: Condition index and reproduction of the Manila clam *Ruditapes philippinarum* in suspended and bottom culture (T. Matsumoto, Y. Ishihi, M. Awaji and J. Higano)
 - 34: Report of a novel flagellated parasite in the Manila clam *Ruditapes philippinarum* on the west coast of Korea (K.I. Park)
 - 35: Geographical distribution of mitochondrial COII haplotypes in the brackish water clam, *Corbicula japonica* (Yamato-Shijimi), around the Japanese archipelago (M. Yamada, R. Ishibashi, K. Toyoda, K. Kawamura and A. Komaru)
 - 36: Effect of different artificial sea waters on survival and growth of adult Manila clam, *Ruditapes philippinarum* (M.N.D. Khan¹, A.M. Shahabuddin¹, N. Arisman, D. Saha, T. Araki and T. Yoshimatsu)
 - 37: Method for detection of colored sand using image analysis and its application to monitoring sediment movement on a tidal flat (A. Kurosawa, T. Okabe, Y. Aoki and S. Kato)
 - 38: Four-year investigations into asari clam and molluscan community in tidal flats after the 2011 Tohoku earthquake (K. Okoshi, W. Shinohara and S. Shimizu)
 - 39: Effect of alien predator *Euspira fortunei* on the distribution and abundance of asari clam after the 2011 Tohoku earthquake and tsunami (M. Suzuki and K. Okoshi)
 - 40: Asari clam lived in new intertidal flat after subsidence of the 2011 Tohoku earthquake (T. Uemura and K. Okoshi)
 - 41: Development of in situ measurement of clearance rate of the Manila clam under

- suspended culture conditions (S. Watanabe, N. Hasegawa and J. Higano)
- 42: Growth and food environment of the Manila clam in suspended culture (Y. Ishihi, S. Watanabe, T. Matsumoto and J. Higano)
- 43: Relationship between abundance of planktonic larvae and benthic juveniles of asari *Ruditapes philippinarum* in eastern Hokkaido, Japan (N. Hasegawa, T. Onituska, S. Ito and H. Abe)
- 44: Energy budget and neuroenzymatic activity in asari, *Ruditapes philippinarum*, following short-term exposure to the toxic alga, *Heterocapsa circularisquama*: implications for reproductive success (L. Basti, S. Nagai, S. Segawa and Y. Tanaka)
- 45: Evaluating the growth and mortality of transplanted asari *Ruditapes philippinarum* juveniles in the Matsusaka region, Mie prefecture, Japan (R. Nambu, Y. Fujioka, N. Hasegawa, T. Kudo and H. Kuwahara)
- 46: Risk assessment on the mortality of asari clam under oxygen deficiency (J. Higano and T. Mizuno)

